



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/12, C07K 14/47, C12N 5/10		A2	(11) International Publication Number: WO 99/55862
			(43) International Publication Date: 4 November 1999 (04.11.99)
(21) International Application Number: PCT/JP99/02226			(81) Designated States: AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 27 April 1999 (27.04.99)			
(30) Priority Data: 10/119395 28 April 1998 (28.04.98) JP			
(71) Applicants (for all designated States except US): SAGAMI CHEMICAL RESEARCH CENTER [JP/JP]; 4-1, Nishi-Ohnuma 4-chome, Sagamihara-shi, Kanagawa 229-0012 (JP). PROTEGENE INC. [JP/JP]; 2-20-3, Naka-cho, Meguro-ku, Tokyo 153-0065 (JP).			
(72) Inventors; and (75) Inventors/Applicants (for US only): KATO, Seishi [JP/JP]; 3-46-50, Wakamatsu, Sagamihara-shi, Kanagawa 229-0014 (JP). KIMURA, Tomoko [JP/JP]; 302, 4-1-28, Nishiikuta, Tama-ku, Kawasaki-shi, Kanagawa 214-0037 (JP).			
(74) Agents: AOYAMA, Tamotsu et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).			Published Without international search report and to be republished upon receipt of that report.
(54) Title: HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAS ENCODING THESE PROTEINS			
(57) Abstract			
<p>A protein comprising any of the amino acid sequences represented by Sequence Nos. 1 to 9, a DNA coding for said protein, exemplified by a cDNA comprising any of the base sequences represented by Sequence Nos. 10 to 18, and an expression vector of said cDNA as well as an eucaryotic cell expressing said cDNA. Said protein and eucaryotic cell having said protein on the membrane surface can be provided by expression of a cDNA coding for a human protein having a transmembrane domain and of a recombinant of the human cDNA.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## DESCRIPTION

HUMAN PROTEINS HAVING TRANSMEMBRANE  
DOMAINS AND DNAs ENCODING THESE PROTEINS

5

TECHNICAL FIELD

The present invention relates to human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells  
10 expressing said cDNAs. The proteins of the present invention can be employed as pharmaceuticals or as antigens for preparing antibodies against said proteins. The human cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources for the gene therapy. Furthermore,  
15 the cDNAs can be utilized as gene sources for large-scale production of the proteins encoded by said cDNAs. Cells, wherein these membrane protein genes are introduced and membrane proteins are expressed in large amounts, can be utilized for detection of the corresponding ligands, screening  
20 of novel low-molecular pharmaceuticals, and so on.

BACKGROUND ART

Membrane proteins play important roles, as signal receptors, ion channels, transporters, etc. in the material  
25 transportation and the information transmission which are mediated by the cell membrane. Examples thereof include receptors for a variety of cytokines, ion channels for the sodium ion, the potassium ion, the chloride ion, etc., transporters for saccharides and amino acids, and so on, where  
30 the genes of many of them have been cloned already.

It has been clarified that abnormalities of these membrane proteins are associated with a number of hitherto-

cryptogenic diseases. For instance, a gene of a membrane protein having twelve transmembrane domains was identified as the gene responsible for cystic fibrosis [Rommens, J. M. et al., Science 245: 1059-1065 (1989)]. In addition, it has been clarified that several membrane proteins act as receptors when a virus infects the cells. For instance, HIV-1 is revealed to infect into the cells through mediation of a membrane protein fusin having a membrane protein on the T-cell membrane, a CD-4 antigen, and seven transmembrane domains [Feng, Y. et al., Science 272: 872-877 (1996)]. Therefore, discovery of a new membrane protein is anticipated to lead to elucidation of the causes of many diseases, so that isolation of a new gene coding for the membrane protein has been desired.

Heretofore, owing to difficulty in the purification, many membrane proteins have been isolated by an approach from the gene side. A general method is the so-called expression cloning which comprises transfection of a cDNA library in eucaryotic cells to express cDNAs and then detection of the cells expressing the target membrane protein on the membrane by an immunological technique using an antibody or a physiological technique on the change in the membrane permeability. However, this method is applicable only to cloning of a gene of a membrane protein with a known function.

In general, membrane proteins possess hydrophobic transmembrane domains inside the proteins, wherein, after synthesis thereof in the ribosome, these domains remain in the phospholipid membrane to be trapped in the membrane. Accordingly, the evidence of the cDNA for encoding the membrane protein is provided by determination of the whole base sequence of a full-length cDNA followed by detection of highly hydrophobic transmembrane domains in the amino acid sequence of the protein encoded by said cDNA.



### DISCLOSURE OF INVENTION

The object of the present invention is to provide novel human proteins having transmembrane domains, DNAs coding for said proteins, and expression vectors of said DNAs as well as transformation eucaryotic cells that are capable of expressing said DNAs.

As the result of intensive studies, the present inventors have been successful in cloning of cDNAs coding for proteins having transmembrane domains from the human full-length cDNA bank, thereby completing the present invention. In other words, the present invention provides human proteins having transmembrane domains, namely proteins containing any of the amino acid sequences represented by Sequence Nos. 1 to 9. Moreover, the present invention provides DNAs coding for the above-mentioned proteins, exemplified by cDNAs containing any of the base sequences represented by Sequence Nos. 10 to 19, 21, 23, 25, 27, 29, 31, 33 and 35, as well as expression vectors that are capable of expressing any of said DNAs by in vitro translation or in eucaryotic cells and transformation eucaryotic cells that are capable of expressing said DNAs and of producing the above-mentioned proteins.

### BRIEF DESCRIPTION OF DRAWINGS

Fig. 1: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02000.

Fig. 2: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02061.

Fig. 3: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by

clone HP02163.

Fig. 4: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02219.

5 Fig. 5: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02256.

10 Fig. 6: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10390.

Fig. 7: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10474.

15 Fig. 8: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10527.

20 Fig. 9: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10528.

#### BEST MODE FOR CARRYING OUT THE INVENTION

25 The proteins of the present invention can be obtained, for example, by a method for isolation from human organs, cell lines, etc., a method for preparation of peptides by the chemical synthesis, or a method for production with the recombinant DNA technology using the DNAs coding for the transmembrane domains of the present invention, wherein the method for obtainment by the recombinant DNA technology is employed preferably. For instance, in vitro expression of the  
30 proteins can be achieved by preparation of an RNA by in vitro transcription from a vector having one of cDNAs of the present invention, followed by in vitro translation using this RNA as a

template. Also, recombination of the translation region into a suitable expression vector by the method known in the art leads to expression of a large amount of the encoded protein by using prokaryotic cells such as *Escherichia coli*, *Bacillus subtilis*, etc., and eucaryotic cells such as yeasts, insect cells, mammalian cells, etc.

In the case in which one of the proteins of the present invention is produced by expressing the DNA by in vitro translation, the protein of the present invention can be produced in vitro, when the translation region of said cDNA is subjected to recombination to a vector having an RNA polymerase promoter, followed by addition to an in vitro translation system such as a rabbit reticulocyte lysate or a wheat germ extract, containing an RNA polymerase corresponding to the promoter. RNA polymerase inhibitors are exemplified by T7, T3, SP6, and the like. The vectors containing these RNA polymerase inhibitors are exemplified by pKA1, pCDM8, pT3/7 18, pT7/3 19, pBluescript II, and so on. Furthermore, a membrane protein of the present invention can be expressed as the form incorporated in the microsome membrane, when a dog pancreas microsome or the like is added into the reaction system.

In the case in which a protein of the present invention is produced by expressing the DNA using a microorganism such as *Escherichia coli* etc., a recombinant expression vector bearing the translation region in the cDNA of the present invention is constructed in an expression vector having an origin, a promoter, a ribosome-binding site, a cDNA-cloning site, a terminator etc., which can be replicated in the microorganism, and, after transformation of the host cells with said expression vector, the thus-obtained transformant is incubated, whereby the protein encoded by said cDNA can be produced on a large scale in the microorganism. In this case, a protein

fragment containing an optional region can be obtained by carrying out the expression with inserting an initiation codon and a termination codon in front of and behind an optional translation region. Alternatively, a fusion protein with another protein can be expressed. Only a protein portion coding for said cDNA can be obtained by cleavage of said fusion protein with a suitable protease. The expression vector for *Escherichia coli* is exemplified by the pUC system, pBluescript II, the pET expression system, the pGEX expression system, and so on.

In the case in which one of the proteins of the present invention is produced by expressing the DNA in eucaryotic cells, the protein of the present invention can be produced as a transmembrane protein on the cell-membrane surface, when the translation region of said cDNA is subjected to recombination to an expression vector for eucaryotic cells that has a promoter, a splicing region, a poly(A) insertion site, etc., followed by introduction into the eucaryotic cells. The expression vector is exemplified by pKA1, pED6dpc2, pCDM8, pSVK3, pMSG, pSVL, pBK-CMV, pBK-RSV, EBV vector, pRS, pYES2, and so on. Examples of eucaryotic cells to be used in general include mammalian culture cells such as simian kidney cells COS7, Chinese hamster ovary cells CHO, etc., budding yeasts, fission yeasts, silkworm cells, *Xenopus laevis* egg cells, and so on, but any eucaryotic cells may be used, provided that they are capable of expressing the present proteins on the membrane surface. The expression vector can be introduced in the eucaryotic cells by methods known in the art such as the electroporation method, the potassium phosphate method, the liposome method, the DEAE-dextran method, and so on.

After one of the proteins of the present invention is expressed in prokaryotic cells or eucaryotic cells, the

objective protein can be isolated from the culture and purified by a combination of separation procedures known in the art. Such examples include treatment with a denaturing agent such as urea or a surface-active agent, sonication, enzymatic digestion, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and so on.

10       The proteins of the present invention include peptide fragments (more than 5 amino acid residues) containing any partial amino acid sequence in the amino acid sequences represented by Sequence Nos. 1 to 9. These peptide fragments can be utilized as antigens for preparation of antibodies.

15       Hereupon, among the proteins of the present invention, those having the signal sequence are secreted in the form of maturation proteins on the surface of the cells, after the signal sequences are removed. Therefore, these maturation proteins shall come within the scope of the present invention.

20       The N-terminal amino acid sequences of the maturation proteins can be easily identified by using the method for the cleavage-site determination in a signal sequence [Japanese Patent Kokai Publication No. 1996-187100]. Furthermore, some membrane proteins undergo the processing on the cell surface to be

25       converted to the secretory forms. Such proteins or peptides in the secretory forms shall come within the scope of the present invention. When sugar chain-binding sites are present in the amino acid sequences, expression in appropriate eucaryotic cells affords proteins wherein sugar chains are added.

30       Accordingly, such proteins or peptides wherein sugar chains are added shall come within the scope of the present invention.

      The DNAs of the present invention include all DNAs coding

for the above-mentioned proteins. Said DNAs can be obtained by using a method by chemical synthesis, a method by cDNA cloning, and so on.

5 The cDNAs of the present invention can be cloned, for example, from cDNA libraries of the human cell origin. These cDNA are synthesized by using as templates poly(A)<sup>+</sup> RNAs extracted from human cells. The human cells may be cells delivered from the human body, for example, by the operation or may be the culture cells. The cDNAs can be synthesized by using  
10 any method selected from the Okayama-Berg method [Okayama, H. and Berg, P., Mol. Cell. Biol. 2: 161-170 (1982)], the Gubler-Hoffman method [Gubler, U. and Hoffman, J. Gene 25: 263-269 (1983)], and so on, but it is preferred to use the capping method [Kato, S. et al., Gene 150: 243-250 (1994)], as  
15 exemplified in Examples, in order to obtain a full-length clone in an effective manner. In addition, commercially available, human cDNA libraries can be utilized. Cloning of the cDNAs of the present invention from the cDNA libraries can be carried out by synthesis of an oligonucleotide on the basis of an  
20 optional portion in the cDNA base sequences of the present invention, followed by screening using this oligonucleotide as the probe according to the colony or plaque hybridization by a method known in the art. In addition, the cDNA fragments of the present invention can be prepared by synthesis of an  
25 oligonucleotide to be hybridized at both termini of the objective cDNA fragment, followed by the usage of this oligonucleotide as the primer for the RT-PCR method from an mRNA isolated from human cells.

30 The cDNAs of the present invention are characterized by containing either of the base sequences represented by Sequence Nos. 10 to 18 or the base sequences represented by Sequence Nos. 19, 21, 23, 25, 27, 29, 31, 33 and 35. Table 1 summarizes the

clone number (HP number), the cells affording the cDNA, the total base number of the cDNA, and the number of the amino acid residues of the encoded protein, for each of the cDNAs.

5

Table 1

Sequence No.	HP No.	Cell	Number of bases	Number of amino acids
1, 10, 19	HP02000	Liver	1705	268
2, 11, 20	HP02061	Saos-2	1759	236
3, 12, 21	HP02163	Saos-2	1069	261
4, 13, 22	HP02219	Stomach Cancer	1759	328
5, 14, 23	HP02256	Stomach Cancer	1697	300
6, 15, 24	HP10390	Stomach Cancer	814	182
7, 16, 25	HP10474	Saos-2	511	66
8, 17, 26	HP10527	Saos-2	1126	183
9, 18, 27	HP10528	Saos-2	2015	324

Hereupon, the same clones as the cDNAs of the present invention can be easily obtained by screening of the cDNA libraries constructed from the human cell lines and human tissues utilized in the present invention by the use of an oligonucleotide probe synthesized on the basis of the cDNA base sequence described in any of Sequence Nos. 10 to 19, 21, 23, 25, 27, 29, 31, 33 and 35.

In general, the polymorphism due to the individual difference is frequently observed in human genes. Accordingly, any cDNA that is subjected to insertion or deletion of one or plural nucleotides and/or substitution with other nucleotides in Sequence Nos. 10 to 19, 21, 23, 25, 27, 29, 31, 33 and 35 shall come within the scope of the present invention.

In a similar manner, any protein that is formed by these modifications comprising insertion or deletion of one or plural amino acids and/or substitution with other amino acids shall

come within the scope of the present invention, as far as the protein possesses the activity of any protein having the amino acid sequences represented by Sequence Nos. 1 to 9.

5 The cDNAs of the present invention include cDNA fragments (more than 10 bp) containing any partial base sequence in the base sequences represented by Sequence Nos. 10 to 18 or in the base sequences represented by Sequence Nos. 19, 21, 23, 25, 27, 29, 31, 33 and 35. Also, DNA fragments consisting of a sense chain and an anti-sense chain shall come within this scope.  
10 These DNA fragments can be utilized as the probes for the gene diagnosis.

In addition to the activities and uses described above, the polynucleotides and proteins of the present invention may exhibit one or more of the uses or biological activities  
15 (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies  
20 or vectors suitable for introduction of DNA).

#### Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for  
25 analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome  
30 markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders;



as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response.

Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions

can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

#### Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one

or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon  $\gamma$ , Schreiber, R.D. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without

limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6-Nordan, R. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

Immune Stimulating or Suppressing Activity

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may

also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a

peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., Science 257:789-792 (1992) and Turka et al., Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function in vivo on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells

that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms.

5 Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking

10 reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases.

15 Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New

20 York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing

25 immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the commoncold, and encephalitis might be alleviated

30 by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be



enhanced in an infected patient by removing T cells from the patient, costimulating the T cells in vitro with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells in vivo.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected ex vivo with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection in vivo.

The presence of the peptide of the present invention

having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which  
5 lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain protein and  $\beta_2$  microglobulin protein or  
10 an MHC class II $\alpha$  chain protein and an MHC class II $\beta$  chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T  
15 cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte  
20 antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among  
25 other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current  
Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing  
30 Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA

78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J.J. and Brunswick, M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that

activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

#### Hematopoiesis Regulating Activity

A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of

factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in  
5 treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity)  
10 useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and  
15 generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell  
20 disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either in-vivo or ex-vivo (i.e., in conjunction with bone marrow  
25 transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

30 Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which

will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lymphohematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

#### Tissue Growth Activity

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the

treatment of burns, incisions and ulcers.

5 A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an  
10 osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair  
15 processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of  
20 bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is  
25 tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in  
30 humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament

tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors ex vivo for return in vivo to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders,



such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

5 Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

10 It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular  
15 endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

20 A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

25 A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

30 Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon);

International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium ).

5 Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

Activin/Inhibin Activity

10 A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are  
15 characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration  
20 of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in  
25 stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such  
30 as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

#### Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays

that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

#### Hemostatic and Thrombolytic Activity

A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

### Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

### Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-

inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

#### Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing,

eliminating or inhibiting factors, agents or cell types which promote tumor growth

Other Activities

5 A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, 10 weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; 15 effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, 20 libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal 25 or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and 30 the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another

material or entity which is cross-reactive with such protein.

#### Examples

The present invention is embodied in more detail by the following examples, but this embodiment is not intended to restrict the present invention. The basic operations and the enzyme reactions with regard to the DNA recombination are carried out according to the literature ["Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Laboratory, 1989]. Unless otherwise stated, restrictive enzymes and a variety of modification enzymes to be used were those available from TAKARA SHUZO. The manufacturer's instructions were used for the buffer compositions as well as for the reaction conditions, in each of the enzyme reactions. The cDNA synthesis was carried out according to the literature [Kato, S. et al., Gene 150: 243-250 (1994)].

##### (1) Preparation of Poly(A)<sup>+</sup> RNA

The osteosarcoma cell line Saos-2 (ATCC HTB 85), tissues of stomach cancer delivered by the operation, and the liver were used for human cells to extract mRNAs. The cell line was incubated by a conventional procedure.

After about 1 g of the human cells was homogenized in 20 ml of a 5.5 M guanidinium thiocyanate solution, a total mRNA was prepared according to the literature [Okayama, H. et al., "Method in Enzymology", Vol. 164, Academic Press, 1987]. This was subjected to chromatography on oligo(dT)-cellulose column washed with a 20 mM Tris-hydrochloride buffer solution (pH 7.6), 0.5 M NaCl, and 1 mM EDTA to obtain a poly(A)<sup>+</sup> RNA according to the above-described literature.

##### (2) Construction of cDNA Library

Ten micrograms of the above-mentioned poly(A)<sup>+</sup> RNA were



dissolved in a 100 mM Tris-hydrochloride buffer solution (pH 8), one unit of an RNase-free, bacterial alkaline phosphatase was added, and the reaction was run at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM sodium acetate (pH 6), 1 mM EDTA, 0.1% 2-mercaptoethanol, and 0.01% Triton X-100. Thereto was added one unit of a tobacco-origin acid pyrophosphatase (Epicentre Technologies) and a total 100  $\mu$ l volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a solution of a decapped poly(A)<sup>+</sup> RNA.

The decapped poly(A)<sup>+</sup> RNA and 3 nmol of a chimeric DNA-RNA oligonucleotide (5'-dG-dG-dG-dG-dA-dA-dT-dT-dC-dG-dA-G-G-A-3') were dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 0.5 mM ATP, 5 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, and 25% polyethylene glycol, whereto was added 50 units of T4RNA ligase and a total 30  $\mu$ l volume of the resulting mixture was reacted at 20°C for 12 hours. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a chimeric-oligo-capped poly(A)<sup>+</sup> RNA.

After digestion of vector pKA1 (Japanese Patent Kokai Publication No. 1992-117292) developed by the present inventors with KpnI, about 60 dT tails were added using a terminal transferase. A vector primer to be used below was prepared by digestion of this product with EcoRV to remove a dT tail at one side.

After 6  $\mu$ g of the previously-prepared chimeric-oligo-capped poly(A)<sup>+</sup> RNA was annealed with 1.2  $\mu$ g of the vector

primer, the resulting product was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 8.3), 75 mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM dithiothreitol, and 1.25 mM dNTP (dATP + dCTP + dGTP + dTTP), 200 units of a reverse transcriptase (GIBCO-BRL) were added, and the reaction in a total 20 µl volume was run at 42°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM NaCl, 10 mM MgCl<sub>2</sub>, and 1 mM dithiothreitol. Thereto were added 100 units of EcoRI and a total 20 µl volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 20 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM KCl, 4 mM MgCl<sub>2</sub>, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, and 50 µg/ml of the bovine serum albumin. Thereto were added 60 units of an *Escherichia coli* DNA ligase and the resulting mixture was reacted at 16°C for 16 hours. To the reaction solution were added 2 µl of 2 mM dNTP, 4 units of *Escherichia coli* DNA polymerase I, and 0.1 unit of *Escherichia coli* RNase H and the resulting mixture was reacted at 12°C for one hour and then at 22°C for one hour.

Next, the cDNA-synthesis reaction solution was used for transformation of *Escherichia coli* DH12S (GIBCO-BRL). The transformation was carried out by the electroporation method. A portion of the transformant was sprayed on the 2xYT agar culture medium containing 100 µg/ml ampicillin and the mixture was incubated at 37°C overnight. A colony formed on the agar medium was picked up at random and inoculated on 2 ml of the 2xYT culture medium containing 100 µg/ml ampicillin. After incubation at 37°C overnight, the culture mixture was

centrifuged to separate the mycelia, from which a plasmid DNA was prepared by the alkaline lysis method. The plasmid DNA was subjected to double digestion with EcoRI and NotI, followed by 0.8% agarose gel electrophoresis, to determine the size of the  
5 cDNA insert. Furthermore, using the thus-obtained plasmid as a template, the sequence reaction was carried out by using an M13 universal primer labeled with a fluorescent dye and a Taq polymerase (a kit of Applied Biosystems) and then the product was examined with a fluorescent DNA sequencer (Applied  
10 Biosystems) to determine an about 400-bp base sequence at the 5'-terminus of the cDNA. The sequence data were filed as the homo/protein cDNA bank database.

(3) Selection of cDNAs Encoding Proteins Having Transmembrane Domains

15 A base sequence registered in the homo/protein cDNA bank was converted to three frames of amino acid sequences and the presence or absence of an open reading frame (ORF) beginning from the initiation codon was examined. Then, the selection was made for the presence of a signal sequence that is  
20 characteristic to a secretory protein at the N-terminus of the portion encoded by the ORF. These clones were sequenced from the both 5' and 3' directions by the use of the deletion method using exonuclease III to determine the whole base sequence. The hydrophobicity/hydrophilicity profiles were obtained for  
25 proteins encoded by the ORF by the Kyte-Doolittle method [Kyte, J. & Doolittle, R. F., J. Mol. Biol. 157: 105-132 (1982)] to examine the presence or absence of a hydrophobic region. In the case in which there is a hydrophobic region of a putative transmembrane domain in the amino acid sequence of an encoded  
30 protein, this protein was judged as a membrane protein.

(4) Functional Verification of Secretory Signal Sequence or Transmembrane Domains

It was verified by the method described in the literature [Yokoyama-Kobayashi, M. et al., Gene 163: 193-196 (1995)] that the N-terminal hydrophobic region in the secretory protein clone candidate obtained in the above-mentioned steps functions  
5 as a secretory signal sequence. First, the plasmid containing the target cDNA was cleaved at an appropriate restriction enzyme site existing at the downstream of the portion expected for encoding the secretory signal sequence. In the case in which this restriction site was a protruding terminus, the site  
10 was blunt-ended by the Klenow treatment or treatment with the T4DNA polymerase. Digestion with HindIII was further carried out and a DNA fragment containing the SV40 promoter and a cDNA encoding the secretory signal sequence at the downstream of the promoter was separated by agarose gel electrophoresis. The  
15 resulting fragment was inserted between HindIII in pSSD3 (DDBJ/EMBL/GenBank Registration No. AB007632) and a restriction enzyme site selected so as to match with the urokinase-coding frame, thereby constructing a vector expressing a fusion protein of the secretory signal sequence of the target cDNA and  
20 the urokinase protease domain.

After *Escherichia coli* (host: JM109) bearing the fusion-protein expression vector was incubated at 37°C for 2 hours in 2 ml of the 2xYT culture medium containing 100 µg/ml of ampicillin, the helper phage M13K07 (50 µl) was added and the  
25 incubation was continued at 37°C overnight. A supernatant separated by centrifugation underwent precipitation with polyethylene glycol to obtain single-stranded phage particles. These particles were suspended in 100 µl of 1 mM Tris-0.1 mM EDTA, pH 8 (TE). Also, there were used as controls suspensions  
30 of single-stranded phage particles prepared in the same manner from pSSD3 and from the vector pKAl-UPA containing a full-length cDNA of urokinase [Yokoyama-Kobayashi, M. et al., Gene

163: 193-196 (1995)].

The culture cells originating from the simian kidney, COS7, were incubated at 37°C in the presence of 5% CO<sub>2</sub> in the Dulbecco's modified Eagle's culture medium (DMEM) containing 10% fetal calf albumin. Into a 6-well plate (Nunc Inc., 3 cm in the well diameter) were inoculated  $1 \times 10^5$  COS7 cells and incubation was carried out at 37°C for 22 hours in the presence of 5% CO<sub>2</sub>. After the culture medium was removed, the cell surface was washed with a phosphate buffer solution and then washed again with DMEM containing 50 mM Tris-hydrochloric acid (pH 7.5) (TDMEM). To the resulting cells was added a suspension of 1 µl of the single-stranded phage suspension, 0.6 ml of the DMEM culture medium, and 3 µl of TRANSFECTAM (IBF Inc.) and the resulting mixture was incubated at 37°C for 3 hours in the presence of 5% CO<sub>2</sub>. After the sample solution was removed, the cell surface was washed with TDMEM, 2 ml per well of DMEM containing 10% fetal calf albumin was added, and the incubation was carried out at 37°C for 2 days in the presence of 5% CO<sub>2</sub>.

To 10 ml of 50 mM phosphate buffer solution (pH 7.4) containing 2% bovine fibrinogen (Miles Inc.), 0.5% agarose, and 1 mM calcium chloride were added 10 units of human thrombin (Mochida Pharmaceutical Co., Ltd.) and the resulting mixture was solidified in a plate of 9 cm in diameter to prepare a fibrin plate. Ten microliters of the culture supernatant of the transfected COS7 cells were spotted on the fibrin plate, which was incubated at 37°C for 15 hours. In the case in which a clear circle appears on the fibrin plate, it is judged that the cDNA fragment codes for the amino acid sequence functioning as a secretory signal sequence. On the other hand, in case in which a clear circle is not formed, the cells were washed well, then the fibrin sheet was placed on the cells, and incubation was carried out at 37°C for 15 hours. In case in which a clear

portion is formed on the fibrin sheet, it indicates that the urokinase activity was expressed on the cell surface. In other words, the cDNA fragment is judged to code for the transmembrane domains.

5 (5) Protein Synthesis by In Vitro Translation

The plasmid vector bearing the cDNA of the present invention was used for in vitro transcription/translation with a T<sub>N</sub>T rabbit reticulocyte lysate kit (Promega). In this case, [<sup>35</sup>S]methionine was added to label the expression product with a radioisotope. Each of the reactions was carried out according to the protocols attached to the kit. Two micrograms of the plasmid was reacted at 30°C for 90 minutes in a total 25 µl volume of the reaction solution containing 12.5 µl of T<sub>N</sub>T rabbit reticulocyte lysate, 0.5 µl of a buffer solution (attached to kit), 2 µl of an amino acid mixture (methionine-free), 2 µl of [<sup>35</sup>S]methionine (Amersham) (0.37 MBq/µl), 0.5 µl of T7RNA polymerase, and 20 U of RNasin. To 3 µl of the resulting reaction solution was added 2 µl of the SDS sampling buffer (125 mM Tris-hydrochloric acid buffer, pH 6.8, 120 mM 2-mercaptoethanol, 2% SDS solution, 0.025% bromophenol blue, and 20% glycerol) and the resulting mixture was heated at 95°C for 3 minutes and then subjected to SDS-polyacrylamide gel electrophoresis. The molecular weight of the translation product was determined by carrying out the autoradiography.

25 (6) Expression by COS7

*Escherichia coli* bearing the expression vector of the protein of the present invention was infected with helper phage M13KO7 and single-stranded phage particles were obtained by the above-mentioned procedure. The thus-obtained phage was used for introducing each expression vector in the culture cells originating from the simian kidney, COS7. After incubation at 37°C for 2 days in the presence of 5% CO<sub>2</sub>, the incubation was

continued for one hour in the culture medium containing [<sup>35</sup>S]cystine or [<sup>35</sup>S]methionine. Collection and dissolution of the cells, followed by subjecting to SDS-PAGE, allowed to observe the presence of a band corresponding to the expression product of each protein, which did not exist in the COS7 cells.

#### (7) Northern Blot Hybridization

Northern blot hybridization was carried out in order to examine the expression pattern in the human tissues. Filters where poly(A)<sup>+</sup> RNAs isolated from each of human tissues are blotted were purchased from Clontech. After excision of a cDNA fragment from the objective clone, followed by agarose-gel electrophoresis to isolate the cDNA fragment, labeling with [<sup>32</sup>P]dCTP (Amersham) was carried out by using a random primer labeling kit (TAKARA SHUZO). The hybridization was carried out by using a solution attached to the blot paper according to the protocol.

#### (8) Clone Examples

<HP02000> (Sequence Nos. 1, 10, and 19)

Determination of the whole base sequence of the cDNA insert of clone HP02000 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a 186-bp 5'-nontranslation region, an 807-bp ORF, and a 712-bp 3'-nontranslation region. The ORF codes for a protein consisting of 268 amino acid residues and there existed two putative transmembrane domains. Figure 1 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 31 kDa that was almost identical with the molecular weight of 30,481 predicted from the ORF. When expressed in COS 7 cells, an expression product of about 32 kDa was observed in the membrane fraction.

10

Table 2

25

RN GAIGQTVLAGLAFTVRNWHHLHLMSVPIFFLLVPTRWLSEARWLIMTNKLQKGLKELI



Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. AA680184) in EST, but any of the sequences was shorter than the present cDNAs and was not found to contain the initiation codon.

An investigation of the expression pattern in the tissues by northern blot hybridization using the cDNA fragment of the present invention has revealed the expression only in the liver.

The rat organic cation transporter has been found as a membrane protein associated with a drug excretion in the kidney [Grundemann, D. et al., Nature 372: 549-552 (1994)]. Accordingly, the protein of the present invention that is its homologue is considered to possess a similar function and can be utilized for the diagnosis and treatment of diseases that are associated with abnormalities of this enzyme. Furthermore, this is considered to be associated with a drug excretion, so that the cells expressing this protein can be used as a tool for designing this drug. In addition, since this protein is expressed specifically in the liver, a substance prepared so as to possess an affinity with this protein can be applied to the drug delivery system to the liver.

<HP02061> (Sequence Nos. 2, 11, and 21)

Determination of the whole base sequence of the cDNA insert of clone HP02061 obtained from cDNA libraries of human osteosarcoma cell line Saos-2 revealed the structure consisting of a 141-bp 5'-nontranslation region, a 711-bp ORF, and a 907-bp 3'-nontranslation region. The ORF codes for a protein consisting of 236 amino acid residues and there existed two putative transmembrane domains. Figure 2 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation

resulted in formation of a translation product of 26 kDa that was almost identical with the molecular weight of 25,593 predicted from the ORF.

The search of the protein data base by using the amino acid sequence of the present protein revealed that the protein was analogous to the human neuroendocrine-specific protein C (PIR Accession No. I60904). Table 3 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the human neuroendocrine-specific protein C (PC). Therein, the marks of -, \*, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively. The C-terminal 187 amino acid residues possessed a homology of 59.9% with the human neuroendocrine-specific protein C.

Table 3

20	HS MAEPSAATQSHSISSSSFGEPSAPGGGGSPGACPALGTKSCSSSCAVHDLIFWRDVKKT <span style="float: right;">**.....*</span>
	PC MQATADSTKMDCVWSNWKSQAIDLLYWRDIKQT HS GFVFGTTLIMLLSLAASFVISVVSYLILALLSVTISFRIYKSVIQAVQKSEEGHPFKAYL <span style="float: right;">*...*. *...*.***.***.*** ** **..*****.*****.*****</span>
25	PC GIVFGSFLLLLFSLTQFSVVSVVAYLALAALSATISFRIYKSVLQAVQKTDEGHPFKAYL HS DVDITLSSEAFHNYMNAAMVHINRALKLIIRLFLVEDLVDSLKLAVFMWLMTYVGAVFNG <span style="float: right;">...**** *...* . ..*...* . *****.*****.***.***.*****.***</span>
	PC ELEITLSQEQIQKYTDCLQFYVNSTLKELRRLFLVQDLVDSLKFVLMWLLTYVGALFNG HS ITLLILAELLIFSVPVIVYEKYKTQIDHYVGIARDQTKSIVEKIQAKLPGIKKKAE <span style="float: right;">***** . *...*.*** *...***.***.***.***.***.*** **..**</span>
30	PC LTLILLMAVVSMTLPVVVVKHQAQIDQYLGLVIRTHINAVVAKIQAKIPG-AKRHAE

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. AA362885) in EST, but, since they are partial  
5 sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

<HP02163> (Sequence Nos. 3, 12, and 23)

Determination of the whole base sequence of the cDNA  
10 insert of clone HP02163 obtained from cDNA libraries of human osteosarcoma cell line Saos-2 revealed the structure consisting of a 179-bp 5'-nontranslation region, a 786-bp ORF, and a 104-bp 3'-nontranslation region. The ORF codes for a protein consisting of 261 amino acid residues and there existed one  
15 putative transmembrane domain. Figure 3 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 30 kDa that was almost identical with the molecular weight of 29,932  
20 predicted from the ORF. When expressed in COS 7 cells, an expression product of about 28 kDa was observed in the membrane fraction.

The search of the protein data base using the amino acid sequence of the present protein has revealed the presence of  
25 sequences that were analogous to a yeast hypothetical protein of 29.4 kDa (SWISS-PROT Accession No. P36039). Table 4 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the yeast hypothetical protein of 29.4 kDa (SC). Therein, the marks of -,  
30 \*, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively.

The both proteins possessed a homology of 33.2% in the entire region.

Table 4

[illegible]

25 Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. Z43161) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

30 <HP02219> (Sequence Nos. 4, 13, and 25)

Determination of the whole base sequence of the cDNA insert of clone HP02219 obtained from cDNA libraries of human

stomach cancer revealed the structure consisting of a 58-bp 5'-nontranslation region, a 987-bp ORF, and a 714-bp 3'-nontranslation region. The ORF codes for a protein consisting of 328 amino acid residues and there existed one putative transmembrane domain. Figure 4 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 39 kDa that was almost identical with the molecular weight of 37,299 predicted from the ORF. When expressed in COS 7 cells, an expression product of about 39 kDa was observed in the membrane fraction.

The search of the protein data base using the amino acid sequence of the present protein has revealed that the protein was analogous to *Alabidopsis thaliana* dTDP-glucose 4-6-dehydratase homologue (PIR Accession No. S58282). Table 5 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the *Alabidopsis thaliana* dTDP-glucose 4-6-dehydratase homologue (AT). Therein, the marks of \* and . represent an amino acid residue identical with the protein of the present invention and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 57.2% in 145 amino acid residues at the C-terminal region.

Table 5

---

HS MVS KALLRLV SAVNRRRMKLLLGIALLAYVASVWGNFVNMSFLLNRSIQENGELKIE  
 5 AT RVVVTGGAGFVGSHLVDRLMARGDTVIVVDNFFTGRKENVMHHFSNPNFEMIRHDVVEPI  
 HS SKIEEMVEPLREKIRDLEKSFTQKYPPVKFLSEKDRKRILITGGAGFVGSHLTDKLMMDG  
 AT LLEVDQIYHLACPASPVHYKFNPKTIKTNVVGTLNMLGLAKRVGARFLLTSTSEVYGD  
 HS HEVTIVVDNFFTGRKRNVEHWIGHENFELINHDVVEPLYIEGVEVRVARIFNTFGPRMHMN  
 10 \*\*\*\*\*.\*\*\*\*\*.\*\*\*\*\*..  
 AT LQHPQVETYWGNVNPIGVRSCYDEGKRTAETLTMDYHRGSNVEVRIARIFNTYGPRMCID  
 HS DGRVVS NFILQALQGEPLTVYGSQSQTRAFQYVSDLVNGLVALMNSNVSSPVNLGNPEEH  
 \*\*\*\*\*.\*\*\*.\*\*\*\*\*.\*.\*\*\*.\*.\*\*\*\*\*.\*.\*\*\*.\*.\*\*\*.\*.  
 AT DGRVVS NFVAQALRKEPLTVYGDGKQTRSFQFVSDLVEGLMRLMEGEHVGPFLGNPGEF  
 15 HS TILEFAQLIKNLVGSSEIQFLSEAQDDPQKRKPDIKKAKMLGWEPVVPLEEGLNKAIH  
 \*.\*.\*.....\*.\*.....\*.\*.\*\*\*\*\*.\*\*\*\*\*.\*.\*\*\*\*\*.\*.\*.\*..  
 AT TMLELAKVVQETIDPNANIEFRPNTEDDPHKRKPDIKAKELLGWEPKVSLRQGLPLMVK  
 HS YFRKELEYQANNQYIPKPKPARIKKGRTRHS  
 \*\*..  
 20 AT DFRQRVFGDQKEGSSAAATTTKT TSA

---

Furthermore, the search of the GenBank using the base  
 sequences of the present cDNA has revealed the registration of  
 25 sequences that possessed a homology of 90% or more (for example,  
 Accession No. U46355) in EST, but, since they are partial  
 sequences, it can not be judged whether or not any of these  
 sequences codes for the same protein as the protein of the  
 present invention.

<HP02256> (Sequence Nos. 5, 14, and 27)

Determination of the whole base sequence of the cDNA insert of clone HP02256 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a 131-bp 5'-nontranslation region, a 903-bp ORF, and a 663-bp 3'-nontranslation region. The ORF codes for a protein consisting of 300 amino acid residues and there existed one transmembrane domain at the N-terminus. Figure 5 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 33 kDa that was almost identical with the molecular weight of 32,943 predicted from the ORF. When expressed in COS cells, an expression product of about 30 kDa was observed in the membrane fraction.

The search of the protein data base using the amino acid sequence of the present protein has revealed that the protein was analogous to the *Caenorhabditis elegans* hypothetical protein T11F9.11 (PID Accession No. 1403260). Table 6 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the *Caenorhabditis elegans* hypothetical protein T11F9.11 (CE). Therein, the marks of -, \*, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively.

The both proteins possessed a homology of 41.7% in the entire region.

Table 6

5	HS MKFLLDIILLPLLVCSLESFVKLFIPK---RRKSVTGEIVLITGAGHGIGRLTAYEFA
	**.. . . . * * * . . . *****. *****. * ***** * ***
	CE MDRALDFVKMVVGTLFFIVLNFFKNFLPNGVLPKRSVEGKKVLITGSGSGIGRLMALEFA
	HS KLKSKLVLDINKHGLEETAACKCKGLGAKVHTFVVDCSNREDIYSSAKKVKAIEIGDVSIL
10	** .....* *** . . . * . . ***** * . . . . . **..*.....*
	CE KLGAEVVIWDVKNKGAEETKNQVVKAGGKASTFVVDLSQYKDIHKVAKETKEAVGDIDIL
	HS VNNAGVVYTSDFATQDPQIEKTFEVNVLAHFWITKAFLPAMTKNNHGHIVTVASAAGHV
	*****. ....* . * . *****. * * . . . *****. * .....***** *
	CE INNAGIVTGKKLFDPCDELMEKTMVNTNALFYTAKNFLPSMLEKDNGHLVTIASMAGKT
15	HS SVPFLLAYCSSKFAAVGFHKTLTDELAALQITGVKTTCLCPNFVNTG-F--IKNPSTSLG
	. . *****. * . * * . . . . * . * . . *****. * * * . . . . . *
	CE GCVGLVDYCASKHGAIGCHDSIAMEILAQKKYGVNTTLVCPFFIDTGMFHGVTTKCPALF
	HS PTLPEEVVNRIMHGILTEQKMIFIPSSIAFLTTLERILPERFLAVLKRKISVKFDAVIG
	*.....* . . . .*****. . . . . . . * . . . *
20	CE PILEANYAVECIVEAILTNRPLLCMPKASYLILALIGLLPIESQVMMADFFGTNESMNDF
	HS YKMQAQ
	CE KGRQKND

25

30

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. H61494) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the



present invention.

<HP10390> (Sequence Nos. 6, 15, and 29)

Determination of the whole base sequence of the cDNA insert of clone HP10390 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a 144-bp 5'-nontranslation region, a 549-bp ORF, and a 121-bp 3'-nontranslation region. The ORF codes for a protein consisting of 182 amino acid residues and possessed one transmembrane domain in the N-terminus. Figure 6 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. Introduction of an expression vector, wherein the HindIII-BstXI (treated with T4RNA polymerase) fragment containing a cDNA portion coding for the N-terminal 50 amino acid residues of the present protein was inserted into the HindIII-SmaI site of pSSD3, into the COS7 cells revealed the urokinase activity on the surface of the cells to indicate that the present protein is the type-II membrane protein. In vitro translation resulted in formation of a translation product of 20 kDa that was almost identical with the molecular weight of 20,639 predicted from the ORF. When expressed in COS cells, an expression product of about 19 kDa was observed in the supernatant fraction and the membrane fraction.

The search of the protein data base using the amino acid sequence of the present protein has not identified any known protein having an analogy. Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. AA315322) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

<HP10474> (Sequence Nos. 7, 16, and 31)

Determination of the whole base sequence of the cDNA insert of clone HP10474 obtained from cDNA libraries of human osteosarcoma cell line Saos-2 revealed the structure consisting of a 22-bp 5'-nontranslation region, a 201-bp ORF, and a 288-bp 3'-nontranslation region. The ORF codes for a protein consisting of 66 amino acid residues and possessed one transmembrane domain at the C-terminus. Figure 7 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 10 kDa that was almost identical with the molecular weight of 7,599 predicted from the ORF.

The search of the protein data base using the amino acid sequence of the present protein has not revealed the presence of any known protein having an analogy. Also, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. H30340) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

<HP10527> (Sequence Nos. 8, 17, and 33)

Determination of the whole base sequence of the cDNA insert of clone HP10527 obtained from cDNA libraries of the human osteosarcoma cell line Saos-2 revealed the structure consisting of a 113-bp 5'-nontranslation region, a 552-bp ORF, and a 461-bp 3'-nontranslation region. The ORF codes for a protein consisting of 183 amino acid residues and possessed three putative transmembrane domains. Figure 8 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. As the result of in

vitro translation, there was produced a translation product of about 21 kDa, which is nearly equal to a molecular weight of 21,111 as expected from ORF.

5 The search of the protein data base using the amino acid sequence of the present protein has not revealed the presence of any known protein having an analogy. Also, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. AA310892) in EST, but, 10 since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

<HP10528> (Sequence Nos. 9, 18, and 35)

Determination of the whole base sequence of the cDNA 15 insert of clone HP10528 obtained from cDNA libraries of the human osteosarcoma cell line Saos-2 revealed the structure consisting of a 53-bp 5'-nontranslation region, a 975-bp ORF, and a 987-bp 3'-nontranslation region. The ORF codes for a protein consisting of 324 amino acid residues and possessed 20 seven putative transmembrane domains. Figure 9 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. As the result of in vitro translation, there was produced a translation product of about 32 kDa, which is nearly equal to a molecular weight of 25 34,227 as expected from ORF.

The search of the protein data base using the amino acid sequence of the present protein has revealed it had an analogy to the epithelial cell growth arrest-inducible gene product (PID Accession No. 998569). Table 7 shows the comparison of the 30 amino acid sequence between the human protein of the present invention (HP) and the epithelial cell growth arrest-inducible gene product (GA). Therein, the marks of -, \*, and . represent

a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 34.7% in the entire region.

5

Table 7

---

	HS	MGPWGEPELLVWRPEAVASEPPVPVGLGVKLGA LVLLLVLTLLCSLVPICVLRPPGANHE
		*..*** * **.* **.*..* .....
10	GA	MEQLLGIKLGCLFALLALTIGCGLTPICFKWFQIDAAR
	HS	GSASRQKALSIVSCFAGGVFLATCLLDLLPDYLA AIDEALAALHV-----
		* . * .*. *..*.....* .....
	GA	GHHRR--VLRLLGCISAGVFLGAGFMHMTAEALEEIESQIQKFMVQNRSASERNSSGDAD
	HS	--TIQFPLQEFILAMGFFLVLMVMEQITLAYKEQSGPSPLEETRALLGTVNGGPQHWHDGP
15		...* *. *..*.....* .....
	GA	SAHMEYPYGELIISLGFFLVFFLESIALQC----CPGA-AGGSTVQDEEWGGAHIF---E
	HS	GVPQASGAPATPSALRACVLVFSIALHSVFEG LAVGLQDRARAMELCLALLLHKGILAV
		... ..* ..*.....* .....
	GA	LHSHGHLPSPSKGPLRALVLLLSLSFHSVFEG LAVGLQPTVAATVQLCLAVLAHKGLVVF
20	HS	SLSRLQLQSHLRAQVVAGCGILFSCMTPLGIGLGAALAES-AGPLHQLAQSVLEGMAAGT
		....**.. ... ..*..*.....* .....
	GA	GVGMLVHLGTSSRWAVFSILLALMSPLGLAVGLAVTGGDSEGGRGLAQAVLEGVAAGT
	HS	FLYITFLEILPQELASSEQRILKVILLAGFALLTG LLIQI
		**.....*.....*.....*
25	GA	ELYVTFLLEILPRELASPEAPLAKWSCVAAGEAFMAFIALWA

---

30

The search of the protein data base using the amino acid sequence of the present protein has not revealed the presence of any known protein having an analogy. Also, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of

90% or more (for example, Accession No. AA206511) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

5

#### INDUSTRIAL APPLICABILITY

The present invention provides human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells  
10 expressing said cDNAs. All of the proteins of the present invention exist in the cell membrane, so that they are considered to be proteins controlling the proliferation and the differentiation of the cells. Accordingly, the proteins of the present invention can be employed as pharmaceuticals such as  
15 carcinostatic agents relating to the control of the proliferation and the differentiation of the cells or as antigens for preparing antibodies against said proteins. The cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources for the gene therapy.  
20 Furthermore, the cDNAs can be utilized for large-scale expression of said proteins. Cells, wherein these membrane protein genes are introduced to possess said proteins on the membrane surface, can be utilized for detection of the corresponding ligands, screening of novel low-molecular  
25 pharmaceuticals, and so on.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide  
30 sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited

to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, Trends Pharmacol. Sci. 15(7): 250-254; Lavarosky et al., 1997, Biochem. Mol. Med. 62(1): 11-22; and Hampel, 1998, Prog. Nucleic Acid Res. Mol. Biol. 58: 1-39; all of which are incorporated by reference herein). Transgenic animals that have multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially

or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, Bioessays 14(9): 629-633; Zwaal et al., 1993, Proc. Natl. Acad. Sci. USA 90(16): 7431-7435; Clark et al., 1994, Proc. Natl. Acad. Sci. USA 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination, preferably detected by positive/negative genetic selection strategies (Mansour et al., 1988, Nature 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms part or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information.

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at

least 25%(more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

Species homologs of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide, as determined by those of skill in the art. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous, or related to that encoded by the polynucleotides.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides



disclosed herein.

The present invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent conditions, and most preferably highlystringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Table

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) <sup>‡</sup>	Hybridization Temperature and Buffer <sup>†</sup>	Wash Temperature and Buffer <sup>†</sup>
A	DNA : DNA	≥50	65°C; 1×SSC -or- 42°C; 1×SSC, 50% formamide	65°C; 0.3×SSC
B	DNA : DNA	<50	T <sub>B</sub> *; 1×SSC	T <sub>B</sub> *; 1×SSC
C	DNA : RNA	≥50	67°C; 1×SSC -or- 45°C; 1×SSC, 50% formamide	67°C; 0.3×SSC
D	DNA : RNA	<50	T <sub>D</sub> *; 1×SSC	T <sub>D</sub> *; 1×SSC
E	RNA : RNA	≥50	70°C; 1×SSC -or- 50°C; 1×SSC, 50% formamide	70°C; 0.3×SSC
F	RNA : RNA	<50	T <sub>F</sub> *; 1×SSC	T <sub>F</sub> *; 1×SSC
G	DNA : DNA	≥50	65°C; 4×SSC -or- 42°C; 4×SSC, 50% formamide	65°C; 1×SSC
H	DNA : DNA	<50	T <sub>H</sub> *; 4×SSC	T <sub>H</sub> *; 4×SSC
I	DNA : RNA	≥50	67°C; 4×SSC -or- 45°C; 4×SSC, 50% formamide	67°C; 1×SSC
J	DNA : RNA	<50	T <sub>J</sub> *; 4×SSC	T <sub>J</sub> *; 4×SSC
K	RNA : RNA	≥50	70°C; 4×SSC -or- 50°C; 4×SSC, 50% formamide	67°C; 1×SSC
L	RNA : RNA	<50	T <sub>L</sub> *; 2×SSC	T <sub>L</sub> *; 2×SSC
M	DNA : DNA	≥50	50°C; 4×SSC -or- 40°C; 6×SSC, 50% formamide	50°C; 2×SSC
N	DNA : DNA	<50	T <sub>N</sub> *; 6×SSC	T <sub>N</sub> *; 6×SSC
O	DNA : RNA	≥50	55°C; 4×SSC -or- 42°C; 6×SSC, 50% formamide	55°C; 2×SSC
P	DNA : RNA	<50	T <sub>P</sub> *; 6×SSC	T <sub>P</sub> *; 6×SSC
Q	RNA : RNA	≥50	60°C; 4×SSC -or- 45°C; 6×SSC, 50% formamide	60°C; 2×SSC
R	RNA : RNA	<50	T <sub>R</sub> *; 4×SSC	T <sub>R</sub> *; 4×SSC

5

‡ : The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid

length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

† : SSPE (1×SSPE is 0.15M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub>, and 1.25mM EDTA, pH7.4) can be substituted for SSC (1×SSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

\*T<sub>B</sub> - T<sub>R</sub> : The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T<sub>m</sub>) of the hybrid, where T<sub>m</sub> is determined according to the following equations. For hybrids less than 18 base pairs in length, T<sub>m</sub>(°C)=2(#of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T<sub>m</sub>(°C)=81.5 + 16.6(log<sub>10</sub>[Na<sup>+</sup>]) + 0.41 (%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na<sup>+</sup>] is the concentration of sodium ions in the hybridization buffer ([Na<sup>+</sup>] for 1×SSC=0.165M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and Current Protocols in Molecular Biology, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25%(more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

## CLAIMS

1. A protein comprising any of the amino acid sequences represented by Sequence Nos. 1 to 9.
- 5 2. A DNA coding for any of the proteins as claimed in Claim 1.
3. A cDNA comprising any of the base sequences represented by Sequence Nos. 10 to 18.
- 10 4. The cDNA as claimed in Claim 3 comprising any of the base sequences represented by Sequence Nos. 19, 21, 23, 25, 27, 29, 31, 33 and 35.
5. An expression vector capable of expressing the DNA as claimed in any of Claim 2 to Claim 4 by in vitro translation or in eucaryotic cells.
- 15 6. A transformation eucaryotic cell capable of expressing the DNA as claimed in any of Claim 2 to Claim 4 and producing the protein as claimed in Claim 1.




Fig. 1




Fig. 2




Fig. 3

4/9




Fig. 4






Fig. 5

6/9




Fig. 6

7/9




Fig. 7

8/9




Fig. 8

9/9




Fig. 9

**Sequence listing**

<110> Sagami Chemical Research Center

5 <120> Human Proteins Having Transmembrane Domains and DNAs Encoding these Proteins

<130> 661099

10 <140>

<141>

<150> JP 10-119395

<151> 1998-04-28

15

<160> 36

<170> Windows 95 (Word 98)

20

<210> 1

<211> 268

<212> PRT

<213> Homo sapiens

25

<400> 1

Met Ala Phe Glu Glu Leu Leu Ser Gln Val Gly Gly Leu Gly Arg Phe

1

5

10

15

Gln Met Leu His Leu Val Phe Ile Leu Pro Ser Leu Met Leu Leu Ile

30

20

25

30

Pro His Ile Leu Leu Glu Asn Phe Ala Ala Ala Ile Pro Gly His Arg

35

40

45

Cys Trp Val His Met Leu Asp Asn Asn Thr Gly Ser Gly Asn Glu Thr

50

55

60

35

Gly Ile Leu Ser Glu Asp Ala Leu Leu Arg Ile Ser Ile Pro Leu Asp

65                      70                      75                      80  
 Ser Asn Leu Arg Pro Glu Lys Cys Arg Arg Phe Val His Pro Gln Trp  
                             85                      90                      95  
 Gln Leu Leu His Leu Asn Gly Thr Ile His Ser Thr Ser Glu Ala Asp  
 5                      100                      105                      110  
 Thr Glu Pro Cys Val Asp Gly Trp Val Tyr Asp Gln Ser Tyr Phe Pro  
                             115                      120                      125  
 Ser Thr Ile Val Thr Lys Trp Asp Leu Val Cys Asp Tyr Gln Ser Leu  
                             130                      135                      140  
 10 Lys Ser Val Val Gln Phe Leu Leu Leu Thr Gly Met Leu Val Gly Gly  
                             145                      150                      155                      160  
 Ile Ile Gly Gly His Val Ser Asp Arg Trp Leu Val Glu Ser Ala Arg  
                             165                      170                      175  
 Trp Leu Ile Ile Thr Asn Lys Leu Asp Glu Gly Leu Lys Ala Leu Arg  
 15                      180                      185                      190  
 Lys Val Ala Arg Thr Asn Gly Ile Lys Asn Ala Glu Glu Thr Leu Asn  
                             195                      200                      205  
 Ile Glu Val Val Arg Ser Thr Met Gln Glu Glu Leu Asp Ala Ala Gln  
                             210                      215                      220  
 20 Thr Lys Thr Thr Val Cys Asp Leu Phe Arg Asn Pro Ser Met Arg Lys  
                             225                      230                      235                      240  
 Arg Ile Cys Ile Leu Val Phe Leu Arg Lys Lys Ile Ser Arg Lys Arg  
                             245                      250                      255  
 His Lys Asn Asp Cys Tyr Thr Lys Val Thr Lys Phe  
 25                      260                      265  
  
 <210> 2  
 <211> 236  
 <212> PRT  
 30 <213> Homo sapiens  
  
 <400> 2  
 Met Ala Glu Pro Ser Ala Ala Thr Gln Ser His Ser Ile Ser Ser Ser  
                             1                      5                      10                      15  
 35 Ser Phe Gly Ala Glu Pro Ser Ala Pro Gly Gly Gly Gly Ser Pro Gly

	20	25	30
	Ala Cys Pro Ala Leu Gly Thr Lys Ser Cys Ser Ser Ser Cys Ala Val		
	35	40	45
	His Asp Leu Ile Phe Trp Arg Asp Val Lys Lys Thr Gly Phe Val Phe		
5	50	55	60
	Gly Thr Thr Leu Ile Met Leu Leu Ser Leu Ala Ala Phe Ser Val Ile		
	65	70	75
	Ser Val Val Ser Tyr Leu Ile Leu Ala Leu Leu Ser Val Thr Ile Ser		
	85	90	95
10	Phe Arg Ile Tyr Lys Ser Val Ile Gln Ala Val Gln Lys Ser Glu Glu		
	100	105	110
	Gly His Pro Phe Lys Ala Tyr Leu Asp Val Asp Ile Thr Leu Ser Ser		
	115	120	125
	Glu Ala Phe His Asn Tyr Met Asn Ala Ala Met Val His Ile Asn Arg		
15	130	135	140
	Ala Leu Lys Leu Ile Ile Arg Leu Phe Leu Val Glu Asp Leu Val Asp		
	145	150	155
	Ser Leu Lys Leu Ala Val Phe Met Trp Leu Met Thr Tyr Val Gly Ala		
	165	170	175
20	Val Phe Asn Gly Ile Thr Leu Leu Ile Leu Ala Glu Leu Leu Ile Phe		
	180	185	190
	Ser Val Pro Ile Val Tyr Glu Lys Tyr Lys Thr Gln Ile Asp His Tyr		
	195	200	205
	Val Gly Ile Ala Arg Asp Gln Thr Lys Ser Ile Val Glu Lys Ile Gln		
25	210	215	220
	Ala Lys Leu Pro Gly Ile Ala Lys Lys Lys Ala Glu		
	225	230	235
	<210> 3		
30	<211> 261		
	<212> PRT		
	<213> Homo sapiens		
	<400> 3		
35	Met Ala Gly Pro Glu Leu Leu Leu Asp Ser Asn Ile Arg Leu Trp Val		



4.

1                    5                    10                    15  
 Val Leu Pro Ile Val Ile Ile Thr Phe Phe Val Gly Met Ile Arg His  
                   20                    25                    30  
 Tyr Val Ser Ile Leu Leu Gln Ser Asp Lys Lys Leu Thr Gln Glu Gln  
 5                    35                    40                    45  
 Val Ser Asp Ser Gln Val Leu Ile Arg Ser Arg Val Leu Arg Glu Asn  
                   50                    55                    60  
 Gly Lys Tyr Ile Pro Lys Gln Ser Phe Leu Thr Arg Lys Tyr Tyr Phe  
                   65                    70                    75                    80  
 10 Asn Asn Pro Glu Asp Gly Phe Phe Lys Lys Thr Lys Arg Lys Val Val  
                   85                    90                    95  
 Pro Pro Ser Pro Met Thr Asp Pro Thr Met Leu Thr Asp Met Met Lys  
                   100                    105                    110  
 Gly Asn Val Thr Asn Val Leu Pro Met Ile Leu Ile Gly Gly Trp Ile  
 15                    115                    120                    125  
 Asn Met Thr Phe Ser Gly Phe Val Thr Thr Lys Val Pro Phe Pro Leu  
                   130                    135                    140  
 Thr Leu Arg Phe Lys Pro Met Leu Gln Gln Gly Ile Glu Leu Leu Thr  
                   145                    150                    155                    160  
 20 Leu Asp Ala Ser Trp Val Ser Ser Ala Ser Trp Tyr Phe Leu Asn Val  
                   165                    170                    175  
 Phe Gly Leu Arg Ser Ile Tyr Ser Leu Ile Leu Gly Gln Asp Asn Ala  
                   180                    185                    190  
 Ala Asp Gln Ser Arg Met Met Gln Glu Gln Met Thr Gly Ala Ala Met  
 25                    195                    200                    205  
 Ala Met Pro Ala Asp Thr Asn Lys Ala Phe Lys Thr Glu Trp Glu Ala  
                   210                    215                    220  
 Leu Glu Leu Thr Asp His Gln Trp Ala Leu Asp Asp Val Glu Glu Glu  
                   225                    230                    235                    240  
 30 Leu Met Ala Lys Asp Leu His Phe Glu Gly Met Phe Lys Lys Glu Leu  
                   245                    250                    255  
 Gln Thr Ser Ile Phe  
                   260  
 35 <210> 4

&lt;211&gt; 328

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

5 &lt;400&gt; 4

Met Val Ser Lys Ala Leu Leu Arg Leu Val Ser Ala Val Asn Arg Arg

1 5 10 15

Arg Met Lys Leu Leu Leu Gly Ile Ala Leu Leu Ala Tyr Val Ala Ser

20 25 30

10 Val Trp Gly Asn Phe Val Asn Met Ser Phe Leu Leu Asn Arg Ser Ile

35 40 45

Gln Glu Asn Gly Glu Leu Lys Ile Glu Ser Lys Ile Glu Glu Met Val

50 55 60

Glu Pro Leu Arg Glu Lys Ile Arg Asp Leu Glu Lys Ser Phe Thr Gln

15 65 70 75 80

Lys Tyr Pro Pro Val Lys Phe Leu Ser Glu Lys Asp Arg Lys Arg Ile

85 90 95

Leu Ile Thr Gly Gly Ala Gly Phe Val Gly Ser His Leu Thr Asp Lys

100 105 110

20 Leu Met Met Asp Gly His Glu Val Thr Val Val Asp Asn Phe Phe Thr

115 120 125

Gly Arg Lys Arg Asn Val Glu His Trp Ile Gly His Glu Asn Phe Glu

130 135 140

Leu Ile Asn His Asp Val Val Glu Pro Leu Tyr Ile Glu Gly Val Glu

25 145 150 155 160

Val Arg Val Ala Arg Ile Phe Asn Thr Phe Gly Pro Arg Met His Met

165 170 175

Asn Asp Gly Arg Val Val Ser Asn Phe Ile Leu Gln Ala Leu Gln Gly

180 185 190

30 Glu Pro Leu Thr Val Tyr Gly Ser Gly Ser Gln Thr Arg Ala Phe Gln

195 200 205

Tyr Val Ser Asp Leu Val Asn Gly Leu Val Ala Leu Met Asn Ser Asn

210 215 220

Val Ser Ser Pro Val Asn Leu Gly Asn Pro Glu Glu His Thr Ile Leu

35 225 230 235 240

Glu Phe Ala Gln Leu Ile Lys Asn Leu Val Gly Ser Gly Ser Glu Ile  
245 250 255  
Gln Phe Leu Ser Glu Ala Gln Asp Asp Pro Gln Lys Arg Lys Pro Asp  
260 265 270  
5 Ile Lys Lys Ala Lys Leu Met Leu Gly Trp Glu Pro Val Val Pro Leu  
275 280 285  
Glu Glu Gly Leu Asn Lys Ala Ile His Tyr Phe Arg Lys Glu Leu Glu  
290 295 300  
Tyr Gln Ala Asn Asn Gln Tyr Ile Pro Lys Pro Lys Pro Ala Arg Ile  
10 305 310 315 320  
Lys Lys Gly Arg Thr Arg His Ser  
325  
  
<210> 5  
15 <211> 300  
<212> PRT  
<213> Homo sapiens  
  
<400> 5  
20 Met Lys Phe Leu Leu Asp Ile Leu Leu Leu Leu Pro Leu Leu Ile Val  
1 5 10 15  
Cys Ser Leu Glu Ser Phe Val Lys Leu Phe Ile Pro Lys Arg Arg Lys  
20 25 30  
Ser Val Thr Gly Glu Ile Val Leu Ile Thr Gly Ala Gly His Gly Ile  
25 35 40 45  
Gly Arg Leu Thr Ala Tyr Glu Phe Ala Lys Leu Lys Ser Lys Leu Val  
50 55 60  
Leu Trp Asp Ile Asn Lys His Gly Leu Glu Glu Thr Ala Ala Lys Cys  
65 70 75 80  
30 Lys Gly Leu Gly Ala Lys Val His Thr Phe Val Val Asp Cys Ser Asn  
85 90 95  
Arg Glu Asp Ile Tyr Ser Ser Ala Lys Lys Val Lys Ala Glu Ile Gly  
100 105 110  
Asp Val Ser Ile Leu Val Asn Asn Ala Gly Val Val Tyr Thr Ser Asp  
35 115 120 125

Leu Phe Ala Thr Gln Asp Pro Gln Ile Glu Lys Thr Phe Glu Val Asn  
 130 135 140  
 Val Leu Ala His Phe Trp Thr Thr Lys Ala Phe Leu Pro Ala Met Thr  
 145 150 155 160  
 5 Lys Asn Asn His Gly His Ile Val Thr Val Ala Ser Ala Ala Gly His  
 165 170 175  
 Val Ser Val Pro Phe Leu Leu Ala Tyr Cys Ser Ser Lys Phe Ala Ala  
 180 185 190  
 Val Gly Phe His Lys Thr Leu Thr Asp Glu Leu Ala Ala Leu Gln Ile  
 10 195 200 205  
 Thr Gly Val Lys Thr Thr Cys Leu Cys Pro Asn Phe Val Asn Thr Gly  
 210 215 220  
 Phe Ile Lys Asn Pro Ser Thr Ser Leu Gly Pro Thr Leu Glu Pro Glu  
 225 230 235 240  
 15 Glu Val Val Asn Arg Leu Met His Gly Ile Leu Thr Glu Gln Lys Met  
 245 250 255  
 Ile Phe Ile Pro Ser Ser Ile Ala Phe Leu Thr Thr Leu Glu Arg Ile  
 260 265 270  
 Leu Pro Glu Arg Phe Leu Ala Val Leu Lys Arg Lys Ile Ser Val Lys  
 20 275 280 285  
 Phe Asp Ala Val Ile Gly Tyr Lys Met Lys Ala Gln  
 290 295 300  
  
 <210> 6  
 25 <211> 182  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 6  
 30 Met Lys Gly Trp Gly Trp Leu Ala Leu Leu Gly Ala Leu Leu Gly  
 1 5 10 15  
 Thr Ala Trp Ala Arg Arg Ser Gln Asp Leu His Cys Gly Ala Cys Arg  
 20 25 30  
 Ala Leu Val Asp Glu Leu Glu Trp Glu Ile Ala Gln Val Asp Pro Lys  
 35 35 40 45

Lys Thr Ile Gln Met Gly Ser Phe Arg Ile Asn Pro Asp Gly Ser Gln  
           50                          55                          60  
 Ser Val Val Glu Val Pro Tyr Ala Arg Ser Glu Ala His Leu Thr Glu  
           65                          70                          75                          80  
 5 Leu Leu Glu Glu Ile Cys Asp Arg Met Lys Glu Tyr Gly Glu Gln Ile  
                                   85                          90                          95  
 Asp Pro Ser Thr His Arg Lys Asn Tyr Val Arg Val Val Gly Arg Asn  
                           100                          105                          110  
 Gly Glu Ser Ser Glu Leu Asp Leu Gln Gly Ile Arg Ile Asp Ser Asp  
 10                          115                          120                          125  
 Ile Ser Gly Thr Leu Lys Phe Ala Cys Glu Ser Ile Val Glu Glu Tyr  
           130                          135                          140  
 Glu Asp Glu Leu Ile Glu Phe Phe Ser Arg Glu Ala Asp Asn Val Lys  
           145                          150                          155                          160  
 15 Asp Lys Leu Cys Ser Lys Arg Thr Asp Leu Cys Asp His Ala Leu His  
                           165                          170                          175  
 Ile Ser His Asp Glu Leu  
                           180  
  
 20 <210> 7  
     <211> 66  
     <212> PRT  
     <213> Homo sapiens  
  
 25 <400> 7  
 Met Glu Val Asp Ala Pro Gly Val Asp Gly Arg Asp Gly Leu Arg Glu  
       1                          5                          10                          15  
 Arg Arg Gly Phe Ser Glu Gly Gly Arg Gln Asn Phe Asp Val Arg Pro  
                           20                          25                          30  
 30 Gln Ser Gly Ala Asn Gly Leu Pro Lys His Ser Tyr Trp Leu Asp Leu  
                           35                          40                          45  
 Trp Leu Phe Ile Leu Phe Asp Val Val Val Phe Leu Phe Val Tyr Phe  
           50                          55                          60  
 Leu Pro  
 35          65

&lt;210&gt; 8

&lt;211&gt; 183

&lt;212&gt; PRT

5 &lt;213&gt; Homo sapiens

&lt;400&gt; 8

Met Ala Ser Arg Ala Gly Pro Arg Ala Ala Gly Thr Asp Gly Ser Asp  
 1 5 10 15  
 10 Phe Gln His Arg Glu Arg Val Ala Met His Tyr Gln Met Ser Val Thr  
 20 25 30  
 Leu Lys Tyr Glu Ile Lys Lys Leu Ile Tyr Val His Leu Val Ile Trp  
 35 40 45  
 Leu Leu Leu Val Ala Lys Met Ser Val Gly His Leu Arg Leu Leu Ser  
 15 50 55 60  
 His Asp Gln Val Ala Met Pro Tyr Gln Trp Glu Tyr Pro Tyr Leu Leu  
 65 70 75 80  
 Ser Ile Leu Pro Ser Leu Leu Gly Leu Leu Ser Phe Pro Arg Asn Asn  
 85 90 95  
 20 Ile Ser Tyr Leu Val Leu Ser Met Ile Ser Met Gly Leu Phe Ser Ile  
 100 105 110  
 Ala Pro Leu Ile Tyr Gly Ser Met Glu Met Phe Pro Ala Ala Gln Gln  
 115 120 125  
 Leu Tyr Arg His Gly Lys Ala Tyr Arg Phe Leu Phe Gly Phe Ser Ala  
 25 130 135 140  
 Val Ser Ile Met Tyr Leu Val Leu Val Leu Ala Val Gln Val His Ala  
 145 150 155 160  
 Trp Gln Leu Tyr Tyr Ser Lys Lys Leu Leu Asp Ser Trp Phe Thr Ser  
 165 170 175  
 30 Thr Gln Glu Lys Lys His Lys  
 180

&lt;210&gt; 9

&lt;211&gt; 324

35 &lt;212&gt; PRT

**<400> 9**

	Met	Gly	Pro	Trp	Gly	Glu	Pro	Glu	Leu	Leu	Val	Trp	Arg	Pro	Glu	Ala	
5	1				5					10						15	
	Val	Ala	Ser	Glu	Pro	Pro	Val	Pro	Val	Gly	Leu	Glu	Val	Lys	Leu	Gly	
				20						25						30	
	Ala	Leu	Val	Leu	Leu	Leu	Val	Leu	Thr	Leu	Leu	Cys	Ser	Leu	Val	Pro	
				35						40						45	
10	Ile	Cys	Val	Leu	Arg	Arg	Pro	Gly	Ala	Asn	His	Glu	Gly	Ser	Ala	Ser	
		50					55					60					
	Arg	Gln	Lys	Ala	Leu	Ser	Leu	Val	Ser	Cys	Phe	Ala	Gly	Gly	Val	Phe	
		65					70					75				80	
	Leu	Ala	Thr	Cys	Leu	Leu	Asp	Leu	Leu	Pro	Asp	Tyr	Leu	Ala	Ala	Ile	
15					85					90						95	
	Asp	Glu	Ala	Leu	Ala	Ala	Leu	His	Val	Thr	Leu	Gln	Phe	Pro	Leu	Gln	
				100						105						110	
	Glu	Phe	Ile	Leu	Ala	Met	Gly	Phe	Phe	Leu	Val	Leu	Val	Met	Glu	Gln	
				115						120						125	
20	Ile	Thr	Leu	Ala	Tyr	Lys	Glu	Gln	Ser	Gly	Pro	Ser	Pro	Leu	Glu	Glu	
		130								135						140	
	Thr	Arg	Ala	Leu	Leu	Gly	Thr	Val	Asn	Gly	Gly	Pro	Gln	His	Trp	His	
		145					150					155				160	
	Asp	Gly	Pro	Gly	Val	Pro	Gln	Ala	Ser	Gly	Ala	Pro	Ala	Thr	Pro	Ser	
25					165						170					175	
	Ala	Leu	Arg	Ala	Cys	Val	Leu	Val	Phe	Ser	Leu	Ala	Leu	His	Ser	Val	
				180						185						190	
	Phe	Glu	Gly	Leu	Ala	Val	Gly	Leu	Gln	Arg	Asp	Arg	Ala	Arg	Ala	Met	
				195						200						205	
30	Glu	Leu	Cys	Leu	Ala	Leu	Leu	Leu	His	Lys	Gly	Ile	Leu	Ala	Val	Ser	
		210								215						220	
	Leu	Ser	Leu	Arg	Leu	Leu	Gln	Ser	His	Leu	Arg	Ala	Gln	Val	Val	Ala	
		225					230					235				240	
	Gly	Cys	Gly	Ile	Leu	Phe	Ser	Cys	Met	Thr	Pro	Leu	Gly	Ile	Gly	Leu	
35					245						250					255	

Gly Ala Ala Leu Ala Glu Ser Ala Gly Pro Leu His Gln Leu Ala Gln  
                             260                            265                            270  
 Ser Val Leu Glu Gly Met Ala Ala Gly Thr Phe Leu Tyr Ile Thr Phe  
                             275                            280                            285  
 5 Leu Glu Ile Leu Pro Gln Glu Leu Ala Ser Ser Glu Gln Arg Ile Leu  
                             290                            295                            300  
 Lys Val Ile Leu Leu Leu Ala Gly Phe Ala Leu Leu Thr Gly Leu Leu  
                             305                            310                            315                            320  
 Phe Ile Gln Ile  
 10  
 <210> 10  
 <211> 804  
 <212> DNA  
 <213> Homo sapiens  
 15  
 <400> 10  
 atggcctttg aggagctctt gagtcaagtt ggaggccttg ggagatttca gatgcttcat 60  
 ctgggttttta ttcttccctc totcatgtta ttaatccctc atatactgct agagaacttt 120  
 gctgcagcca ttctgtgtca tcgttgctgg gtccacatgc tggacaataa tactggatct 180  
 20 ggtaatgaaa ctggaatcct cagtgaagat gccctcttga gaatctctat ccactagac 240  
 tcaaactctga ggccagagaa gtgtcgtcgc tttgtccatc ccagtgaggca gcttcttcac 300  
 ctgaatggga ctatccacag cacaagtga gacagacacag aaccctgtgt ggatggctgg 360  
 gtatatgatc aaagctactt cccttcgacc attgtgacta agtgggacct ggtatgtgat 420  
 tatcagtcac tgaaatcagt ggttcaattc ctacttctga ctggaatgct ggtgggaggc 480  
 25 atcataggtg gccatgtctc agacaggtgg ctggtggaat ctgctcgggtg gttgataatc 540  
 accaataaac tagatgaggg cttaaaggca cttagaaaag ttgcacgcac aaatggaata 600  
 aagaatgctg aagaaaccct gaacatagag gttgtaagat ccaccatgca ggaggagctg 660  
 gatgcagcac agacaaaaac tactgtgtgt gacttgttcc gcaaccccag tatgcgtaaa 720  
 aggatctgta tcctgggtatt tttagaaaa aaaatctcaa ggaaaaggca taaaaatgat 780  
 30 tgctacacaa aagtgaccaa attt 804  
 <210> 11  
 <211> 708  
 <212> DNA  
 35 <213> Homo sapiens



&lt;400&gt; 11

atggcggagc cgtcggcggc cactcagtc cattccatct cctcgtcgtc cttcggagcc 60  
 gagccgtccg cgccccggcg cggcgggagc ccaggagcct gccccgccct ggggacgaag 120  
 5 agctgcagct cctcctgtgc ggtgcacgat ctgattttct ggagagatgt gaagaagact 180  
 gggtttgtct ttggcaccac gctgatcatg ctgctttccc tggcagcttt cagtgtcatc 240  
 agtgtggttt cttacctcat cctggctctt ctctctgtca ccatcagctt caggatctac 300  
 aagtccgtca tccaagctgt acagaagtca gaagaaggcc atccattcaa agcctacctg 360  
 gacgtagaca ttactctgtc ctcagaagct ttccataatt acatgaatgc tgccatggtg 420  
 10 cacatcaaca gggccctgaa actcattatt cgtctctttc tggtagaaga tctggttgac 480  
 tccttgaagc tggctgtctt catgtggctg atgacctatg ttggtgctgt ttttaacgga 540  
 atcacccttc taattcttgc tgaactgtc attttcagt tcccagattgt ctatgagaag 600  
 tacaagacct agattgatca ctatgttggc atcgcccgag atcagaccaa gtcaattgtt 660  
 gaaaagatcc aagcaaaact ccctggaatc gccaaaaaaa aggcagaa 708

15

&lt;210&gt; 12

&lt;211&gt; 783

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

20

&lt;400&gt; 12

atggcagggc cagaactgtt gctcgactcc aacatccgcc tctgggtggt cctacccatc 60  
 gttatcatca ctttcttctt aggcattgat cgccactacg tgtccatcct gctgcagagc 120  
 gacaagaagc tcaccacagga acaagtatct gacagtcaag tcctaattcg aagcagagtc 180  
 25 ctcagggaaa atggaaaata cattcccaaa cagtctttct tgacacgaaa atattatttc 240  
 aacaacccag aggatggatt tttcaaaaaa actaaacgga aggtagtgcc accttctcct 300  
 atgactgatc ctactatgtt gacagacatg atgaaaggga atgtaacaaa tgcctccct 360  
 atgattctta ttggtggatg gatcaacatg acattctcag gctttgtcac aaccaaggtc 420  
 ccatttccac tgacctccg ttttaagcct atgttacagc aaggaatcga gctactcaca 480  
 30 ttagatgcat cctgggtgag ttctgcatcc tggacttcc tcaatgtatt tgggcttcgg 540  
 agcatttact ctctgattct gggccaagat aatgccgctg accaatcacg aatgatgcag 600  
 gagcagatga cgggagcagc catggccatg cccgcagaca caaacaagc tttcaagaca 660  
 gagtgggaag ctttgagct gacggatcac cagtgggcac tagatgatgt cgaagaagag 720  
 ctcatggcca aagacctcca cttcgaaggc atgttcaaaa aggaattaca gacctctatt 780  
 35 ttt 783

&lt;210&gt; 13

&lt;211&gt; 984

&lt;212&gt; DNA

5 &lt;213&gt; Homo sapiens

&lt;400&gt; 13

atggtgagca aggcgctgct gcgcctcgtg tctgccgtca accgcaggag gatgaagctg 60  
ctgctgggca tcgccttget ggcctacgtc gcctctgttt ggggcaactt cgtaatatg 120  
10 agctttctac tcaacaggtc tatccaggaa aatggtgaac taaaaattga aagcaagatt 180  
gaagagatgg ttgaaccact aagagagaaa atcagagatt tagaaaaaag ctttaccag 240  
aaatacccac cagtaaagtt tttatcagaa aaggatcgga aaagaatttt gataacagga 300  
ggcgcaggggt tcgtgggctc ccatctaact gacaaactca tgatggacgg ccacgaggtg 360  
accgtggtgg acaatttctt cacgggcagg aagagaaacg tggagcactg gatcggacat 420  
15 gagaacttcg agttgattaa ccacgacgtg gtggagcccc tctacatcga gggcgtggaa 480  
gtgcgagtggt ccagaatctt caacaccttt gggccacgca tgcacatgaa cgatgggcga 540  
gtagtcagca acttcatcct gcaggcgctc cagggggagc cactcacggt atacggatcc 600  
gggtctcaga caagggcggt ccagtacgtc agcgatctag tgaatggcct cgtggctctc 660  
atgaacagca acgtcagcag cccggtcaac ctggggaacc cagaagaaca cacaatccta 720  
20 gaatttgctc agttaattaa aaaccttggt ggtagcggaa gtgaaattca gtttctctcc 780  
gaagcccagg atgaccaca gaaaagaaaa ccagacatca aaaaagcaaa gctgatgctg 840  
gggtgggagc ccgtgggtccc gctggaggaa ggtttaaaca aagcaattca ctacttccgt 900  
aaagaactcg agtaccaggc aaataatcag tacatcccca aaccaaagcc tgccagaata 960  
aagaaaggac ggactcgcca cagc 984

25

&lt;210&gt; 14

&lt;211&gt; 900

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

30

&lt;400&gt; 14

atgaaatttc ttctggacat cctcctgctt ctcccgttac tgatcgtctg ctccctagag 60  
tccttcgtga agctttttat tcctaagagg agaaaatcag tcaccggcga aatcgtgctg 120  
attacaggag ctgggcatgg aattgggaga ctgactgcct atgaatttgc taaacttaaa 180  
35 agcaagctgg ttctctggga tataaataag catggactgg aggaaacagc tgccaaatgc 240

aagggactgg gtgccaaggt tcataccttt gtggttagact gcagcaaccg agaagatatt 300  
 tacagctctg caaagaaggt gaaggcagaa attggagatg ttagtatttt agtaaataat 360  
 gctggtgtag tctatacatc agatttggtt gctacacaag atcctcagat tgaaaagact 420  
 tttgaagtta atgtacttgc acatttctgg actacaaagg catttcttcc tgcaatgacg 480  
 5 aagaataacc atggccatat tgtcactgtg gcttcggcag ctggacatgt ctcggtcccc 540  
 ttcttactgg cttactgttc aagcaagttt gctgctgttg gatttcataa aactttgaca 600  
 gatgaactgg ctgccttaca aataactgga gtcaaaacaa catgtctgtg tcctaatttc 660  
 gtaaactg gcttcatcaa aaatccaagt acaagtttg gacccactct ggaacctgag 720  
 gaagtggtaa acaggctgat gcatgggatt ctgactgagc agaagatgat ttttattcca 780  
 10 tcttctatag cttttttaac aacattggaa aggatccttc ctgagcggtt cctggcagtt 840  
 ttaaaacgaa aaatcagtgt taagtttgat gcagttattg gatataaaat gaaagcgcaa 900

<210> 15

<211> 546

15 <212> DNA

<213> Homo sapiens

<400> 15

atgaaaggct ggggttggtt ggccctgctt ctggggggccc tgctgggaac cgcctgggct 60  
 20 cggaggagcc aggatctcca ctgtggagca tgcagggtc tggaggatga actagaatgg 120  
 gaaattgccc aggtggaccc caagaagacc attcagatgg gatctttccg gatcaatcca 180  
 gatggcagcc agtcagtggg ggaggtgcct tatgcccgct cagaggccca cctcacagag 240  
 ctgctggagg agatatgtga ccggatgaag gagtatggg aacagattga tccttcacc 300  
 catcgcaaga actacgtacg tgtagtgggc cggaatggag aatccagtga actggacct 360  
 25 caaggcatcc gaatcgactc agatattagc ggcaccctca agtttgctg tgagagcatt 420  
 gtggaggaat acgaggatga actcattgaa ttcttttccc gagaggctga caatgttaaa 480  
 gacaaacttt gcagtaagcg aacagatctt tgtgaccatg ccctgcacat atcgcatgat 540  
 gagcta

30 <210> 16

<211> 198

<212> DNA

<213> Homo sapiens

35 <400> 16

15.

	atggaggtgg acgcaccggg tgttgatggt cgagatggtc tccgggagcg gcgaggtttt	60
	agcgagggag ggaggcagaa cttcgatgtg aggcctcagt ctggggcaaa tgggttccc	120
	aaacactcct actggttga cctctggctt ttcactctt tcgatgtggt ggtgtttctc	180
	tttgtgtatt ttttgcca	198
5	<210> 17	
	<211> 549	
	<212> DNA	
	<213> Homo sapiens	
10	<400> 17	
	atggcgtctc gagcaggccc gcgagcggcc ggcaccgacg gcagcgactt tcagcaccgg	60
	gagcgcgtcg ccatgcacta ccagatgagt gtgacctca agtatgaaat caagaagctg	120
	atctacgtac atctggatcat atggctgctg ctggttgcta agatgagcgt gggacacctg	180
15	aggctcttgt cacatgatca ggtggccatg ccctatcagt gggaataccc gtatttgcgtg	240
	agcattttgc cctctctctt gggccttctc tcctttcccc gcaacaacat tagctacctg	300
	gtgctctcca tgatcagcat gggactcttt tccatcgctc cactcattta tggcagcatg	360
	gagatgttcc ctgctgcaca gcagctctac cgccatggca aggcctaccg tttcctcttt	420
	ggtttttctg ccgtttccat catgtacctg gtgttggtgt tggcagtgca agtgcagccc	480
20	tggcagttgt actacagcaa gaagctccta gactcttggt tcaccagcac acaggagaag	540
	aagcataaa	549
	<210> 18	
	<211> 972	
25	<212> DNA	
	<213> Homo sapiens	
	<400> 18	
	atggggccct ggggagagcc agagctcctg gtgtggcgcc ccgaggcggg agcttcagag	60
30	cctccagtgct ctgtggggct ggaggtgaag ttgggggccc tgggtgctgct gctggtgctc	120
	accctcctct gcagcctggt gccatctgt gtgctgcgcc ggccaggagc taacctgaa	180
	ggctcagctt cccgccagaa agccctgagc ctagtaagct gtttcgcggg gggcgtcttt	240
	ttggccaact gtctcctgga cctgctgcct gactacctgg ctgccataga tgaggccctg	300
	gcagccttgc acgtgacgct ccagttccca ctgcaagagt tcactcctggc catgggcttc	360
35	ttcctggtcc tgggtgatgga gcagatcaca ctggcttaca aggagcagtc agggccgtca	420

	cctctggagg aaacaagggc tctgctggga acagtgaatg gtgggcccga gcattggcat	480
	gatgggccag gggccccaca ggcgagtgga gcccagcaa cccctcagc cttgcgtgcc	540
	tgtgtactgg tgttctccct ggccctccac tccgtgttcg aggggctggc ggtagggctg	600
	cagcgagacc gggctcgggc catggagctg tgcctggctt tgctgctcca caagggcatc	660
5	ctggctgtca gcctgtccct gcggctgttg cagagccacc ttagggcaca ggtggtggct	720
	ggctgtggga tcctcttctc atgcatgaca cctctaggca tcgggctggg tgcagctctg	780
	gcagagtcgg caggacctct gcaccagctg gccagctctg tgctagaggg catggcagct	840
	ggcaccttct tctatatcac ctttctggaa atcctgcccc aggagctggc cagttctgag	900
	caaaggatcc tcaaggtcat tctgctccta gcaggctttg cctgctcac tggcctgctc	960
10	ttcatccaaa tc	972
	<210> 19	
	<211> 1705	
	<212> DNA	
15	<213> Homo sapiens	
	<400> 19	
	aagaactgag gaagctcttt ccactacggc tgtattgcac tggtagtcc gggcccatgg	60
	atgagaaatt gatgcgagga tcaatacaag cttaatttga attaataaaa ggaaatattt	120
20	tctccctttg aacttatctc cgtaaagcca ttgtgcctcc tcttgggggt cacgtgttca	180
	caatca atg gcc ttt gag gag ctc ttg agt caa gtt gga ggc ctt ggg	228
	Met Ala Phe Glu Glu Leu Leu Ser Gln Val Gly Gly Leu Gly	
	1 5 10	
	aga ttt cag atg ctt cat ctg gtt ttt att ctt ccc tct ctc atg tta	276
25	Arg Phe Gln Met Leu His Leu Val Phe Ile Leu Pro Ser Leu Met Leu	
	15 20 25 30	
	tta atc cct cat ata ctg cta gag aac ttt gct gca gcc att cct ggt	324
	Leu Ile Pro His Ile Leu Leu Glu Asn Phe Ala Ala Ala Ile Pro Gly	
	35 40 45	
30	cat cgt tgc tgg gtc cac atg ctg gac aat aat act gga tct ggt aat	372
	His Arg Cys Trp Val His Met Leu Asp Asn Asn Thr Gly Ser Gly Asn	
	50 55 60	
	gaa act gga atc ctc agt gaa gat gcc ctc ttg aga atc tct atc cca	420
	Glu Thr Gly Ile Leu Ser Glu Asp Ala Leu Leu Arg Ile Ser Ile Pro	
35	65 70 75	

	cta gac tca aat ctg agg cca gag aag tgt cgt cgc ttt gtc cat ccc	468
	Leu Asp Ser Asn Leu Arg Pro Glu Lys Cys Arg Arg Phe Val His Pro	
	80 85 90	
	cag tgg cag ctt ctt cac ctg aat ggg act atc cac agc aca agt gag	516
5	Gln Trp Gln Leu Leu His Leu Asn Gly Thr Ile His Ser Thr Ser Glu	
	95 100 105 110	
	gca gac aca gaa ccc tgt gtg gat ggc tgg gta tat gat caa agc tac	564
	Ala Asp Thr Glu Pro Cys Val Asp Gly Trp Val Tyr Asp Gln Ser Tyr	
	115 120 125	
10	ttc cct tcg acc att gtg act aag tgg gac ctg gta tgt gat tat cag	612
	Phe Pro Ser Thr Ile Val Thr Lys Trp Asp Leu Val Cys Asp Tyr Gln	
	130 135 140	
	tca ctg aaa tca gtg gtt caa ttc cta ctt ctg act gga atg ctg gtg	660
	Ser Leu Lys Ser Val Val Gln Phe Leu Leu Leu Thr Gly Met Leu Val	
15	145 150 155	
	gga ggc atc ata ggt ggc cat gtc tca gac agg tgg ctg gtg gaa tct	708
	Gly Gly Ile Ile Gly Gly His Val Ser Asp Arg Trp Leu Val Glu Ser	
	160 165 170	
	gct cgg tgg ttg ata atc acc aat aaa cta gat gag ggc tta aag gca	756
20	Ala Arg Trp Leu Ile Ile Thr Asn Lys Leu Asp Glu Gly Leu Lys Ala	
	175 180 185 190	
	ctt aga aaa gtt gca cgc aca aat gga ata aag aat gct gaa gaa acc	804
	Leu Arg Lys Val Ala Arg Thr Asn Gly Ile Lys Asn Ala Glu Glu Thr	
	195 200 205	
25	ctg aac ata gag gtt gta aga tcc acc atg cag gag gag ctg gat gca	852
	Leu Asn Ile Glu Val Val Arg Ser Thr Met Gln Glu Glu Leu Asp Ala	
	210 215 220	
	gca cag acc aaa act act gtg tgt gac ttg ttc cgc aac ccc agt atg	900
	Ala Gln Thr Lys Thr Thr Val Cys Asp Leu Phe Arg Asn Pro Ser Met	
30	225 230 235	
	cgt aaa agg atc tgt atc ctg gta ttt ttg aga aaa aaa atc tca agg	948
	Arg Lys Arg Ile Cys Ile Leu Val Phe Leu Arg Lys Lys Ile Ser Arg	
	240 245 250	
	aaa agg cat aaa aat gat tgc tac aca aaa gtg acc aaa ttt taagaagcct	1000
35	Lys Arg His Lys Asn Asp Cys Tyr Thr Lys Val Thr Lys Phe	

	255	260	265	
	tcatgagctg attggtgggg aaattcagaa aaaaaaatac aggaaaagaa cacaccagaa	1060		
	gggttttttt ccctacaacc agcaagaaca tatattagat acatgaatct caattataat	1120		
	tatggcatta atttgcattt tatttcaaaa ttaacttggtg gggacatgta atctcttgag	1180		
5	caatctgata tttttgggaa gtcctttaaa aagttacaaa tttatcaata aattactagt	1240		
	agataagatg attcagaaac aaaagaaaat cacagaatta ggatgtggct ggctggtgta	1300		
	tgaagcacca tgtgatgaat tcataaagtt gcaaaagtca aaacaatact gtacatgcaa	1360		
	ccagaaatca aaataaatcc agaaatagag acctatataa atgcatttaa tacatgatac	1420		
	ttttgacata ataagccatt ggaaaacgga aagattagat actaaataac attgactatc	1480		
10	tctttgtaaa tacagtcact aaatgatgtt agttactttt ccatggtgga attttaatta	1540		
	ctttttcttt gtaatttttc tctctgtata ttttaaacia atagctggta tagtttacia	1600		
	tattataaag atattgttca aattgaaggg caaaggccag gttcagcaat tttcaaactg	1660		
	tatgtacatt taataaaata actataaatt aaaaaattat atttc	1705		
15	<210> 20			
	<211> 268			
	<212> PRT			
	<213> Homo sapiens			
20	<400> 20			
	Met Ala Phe Glu Glu Leu Leu Ser Gln Val Gly Gly Leu Gly			
	1 5 10			
	Arg Phe Gln Met Leu His Leu Val Phe Ile Leu Pro Ser Leu Met Leu			
	15 20 25 30			
25	Leu Ile Pro His Ile Leu Leu Glu Asn Phe Ala Ala Ala Ile Pro Gly			
	35 40 45			
	His Arg Cys Trp Val His Met Leu Asp Asn Asn Thr Gly Ser Gly Asn			
	50 55 60			
	Glu Thr Gly Ile Leu Ser Glu Asp Ala Leu Leu Arg Ile Ser Ile Pro			
30	65 70 75			
	Leu Asp Ser Asn Leu Arg Pro Glu Lys Cys Arg Arg Phe Val His Pro			
	80 85 90			
	Gln Trp Gln Leu Leu His Leu Asn Gly Thr Ile His Ser Thr Ser Glu			
	95 100 105 110			
35	Ala Asp Thr Glu Pro Cys Val Asp Gly Trp Val Tyr Asp Gln Ser Tyr			

19.

115 120 125  
 Phe Pro Ser Thr Ile Val Thr Lys Trp Asp Leu Val Cys Asp Tyr Gln  
 130 135 140  
 Ser Leu Lys Ser Val Val Gln Phe Leu Leu Leu Thr Gly Met Leu Val  
 5 145 150 155  
 Gly Gly Ile Ile Gly Gly His Val Ser Asp Arg Trp Leu Val Glu Ser  
 160 165 170  
 Ala Arg Trp Leu Ile Ile Thr Asn Lys Leu Asp Glu Gly Leu Lys Ala  
 175 180 185 190  
 10 Leu Arg Lys Val Ala Arg Thr Asn Gly Ile Lys Asn Ala Glu Glu Thr  
 195 200 205  
 Leu Asn Ile Glu Val Val Arg Ser Thr Met Gln Glu Glu Leu Asp Ala  
 210 215 220  
 Ala Gln Thr Lys Thr Thr Val Cys Asp Leu Phe Arg Asn Pro Ser Met  
 15 225 230 235  
 Arg Lys Arg Ile Cys Ile Leu Val Phe Leu Arg Lys Lys Ile Ser Arg  
 240 245 250  
 Lys Arg His Lys Asn Asp Cys Tyr Thr Lys Val Thr Lys Phe  
 255 260 265  
 20  
 <210> 21  
 <211> 1759  
 <212> DNA  
 <213> Homo sapiens  
 25  
 <400> 21  
 agtcagtcctg tcggagtcctg tcctcggagc aggcggagta aagggacttg agcgagccag 60  
 ttgccggatt attctatttc cctccctct ctcccgcccc gtatctcttt tcacccttct 120  
 cccaccctcg ctgcgtagc c atg gcg gag ccg tcg gcg gcc act cag tcc 171  
 30 Met Ala Glu Pro Ser Ala Ala Thr Gln Ser  
 1 5 10  
 cat tcc atc tcc tcg tcg tcc ttc gga gcc gag ccg tcc gcg ccc ggc 219  
 His Ser Ile Ser Ser Ser Ser Phe Gly Ala Glu Pro Ser Ala Pro Gly  
 15 20 25  
 35 ggc ggc ggg agc cca gga gcc tgc ccc gcc ctg ggg acg aag agc tgc 267



	Gly Gly Gly Ser Pro Gly Ala Cys Pro Ala Leu Gly Thr Lys Ser Cys	
	30 35 40	
	agc tcc tcc tgt gcg gtg cac gat ctg att ttc tgg aga gat gtg aag	315
	Ser Ser Ser Cys Ala Val His Asp Leu Ile Phe Trp Arg Asp Val Lys	
5	45 50 55	
	aag act ggg ttt gtc ttt ggc acc acg ctg atc atg ctg ctt tcc ctg	363
	Lys Thr Gly Phe Val Phe Gly Thr Thr Leu Ile Met Leu Leu Ser Leu	
	60 65 70	
	gca gct ttc agt gtc atc agt gtg gtt tct tac ctc atc ctg gct ctt	411
10	Ala Ala Phe Ser Val Ile Ser Val Val Ser Tyr Leu Ile Leu Ala Leu	
	75 80 85 90	
	ctc tct gtc acc atc agc ttc agg atc tac aag tcc gtc atc caa gct	459
	Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr Lys Ser Val Ile Gln Ala	
	95 100 105	
15	gta cag aag tca gaa gaa ggc cat cca ttc aaa gcc tac ctg gac gta	507
	Val Gln Lys Ser Glu Glu Gly His Pro Phe Lys Ala Tyr Leu Asp Val	
	110 115 120	
	gac att act ctg tcc tca gaa gct ttc cat aat tac atg aat gct gcc	555
	Asp Ile Thr Leu Ser Ser Glu Ala Phe His Asn Tyr Met Asn Ala Ala	
20	125 130 135	
	atg gtg cac atc aac agg gcc ctg aaa ctc att att cgt ctc ttt ctg	603
	Met Val His Ile Asn Arg Ala Leu Lys Leu Ile Ile Arg Leu Phe Leu	
	140 145 150	
	gta gaa gat ctg gtt gac tcc ttg aag ctg gct gtc ttc atg tgg ctg	651
25	Val Glu Asp Leu Val Asp Ser Leu Lys Leu Ala Val Phe Met Trp Leu	
	155 160 165 170	
	atg acc tat gtt ggt gct gtt ttt aac gga atc acc ctt cta att ctt	699
	Met Thr Tyr Val Gly Ala Val Phe Asn Gly Ile Thr Leu Leu Ile Leu	
	175 180 185	
30	gct gaa ctg ctc att ttc agt gtc ccg att gtc tat gag aag tac aag	747
	Ala Glu Leu Leu Ile Phe Ser Val Pro Ile Val Tyr Glu Lys Tyr Lys	
	190 195 200	
	acc cag att gat cac tat gtt ggc atc gcc cga gat cag acc aag tca	795
	Thr Gln Ile Asp His Tyr Val Gly Ile Ala Arg Asp Gln Thr Lys Ser	
35	205 210 215	

	att gtt gaa aag atc caa gca aaa ctc cct gga atc gcc aaa aaa aag	843
	Ile Val Glu Lys Ile Gln Ala Lys Leu Pro Gly Ile Ala Lys Lys Lys	
	220 225 230	
	gca gaa taagtacatg gaaaccagaa atgcaacagt tactaaaaca ccatttaata g	900
5	Ala Glu	
	235	
	ttataacgtc gttacttgta ctatgaagga aaatactcag tgtcagcttg agcctgcatt	960
	ccaagctttt tttttaattt ggtgttttct cccatccttt ccctttaacc ctgagtatca	1020
	agcacaaaaa ttgatggact gataaaagaa ctatcttaga actcagaaga agaaagaatc	1080
10	aaattcatag gataagtcaa taccttaatg gtggtagagc ctttacctgt agcttgaaag	1140
	gggaaagatt ggaggtaga gagaaaatga aagaacacct ctgggtcctt ctgtccagtt	1200
	ttcagcacta gtcttactca gctatccatt atagttttgc ccttaagaag tcatgattaa	1260
	cttatgaaaa aattatttgg ggacaggagt gtgatacctt ccttggtttt tttttgcagc	1320
	cctcaaatcc tatcttctcg cccacaatg tgagcagcta cccctgatac tccttttctt	1380
15	taatgattta actatcaact tgataaataa cttatagggtg atagtataa ttcttgattc	1440
	caagaatgcc atctgataaa aaagaataga aatggaaagt gggactgaga gggagtcagc	1500
	aggcatgctg cgggtggcggg cactccctct gccactatcc ccaggggaagg aaaggctccg	1560
	ccatttgagg aagtgggttc tacgtcactg gacaccgggt ctgagcatta gtttgagaac	1620
	tcgttccga atgtgctttc ctccctctcc cctgccacc tcaagtttaa taaataaggt	1680
20	tgtacttttc ttactataaa ataaatgtct gtaactgctg tgcactgctg taaacttggt	1740
	agagaaaaaa ataacctgc	1759
	<210> 22	
	<211> 236	
25	<212> PRT	
	<213> Homo sapiens	
	<400> 22	
	Met Ala Glu Pro Ser Ala Ala Thr Gln Ser	
30	1 5 10	
	His Ser Ile Ser Ser Ser Ser Phe Gly Ala Glu Pro Ser Ala Pro Gly	
	15 20 25	
	Gly Gly Gly Ser Pro Gly Ala Cys Pro Ala Leu Gly Thr Lys Ser Cys	
	30 35 40	
35	Ser Ser Ser Cys Ala Val His Asp Leu Ile Phe Trp Arg Asp Val Lys	

45 50 55  
 Lys Thr Gly Phe Val Phe Gly Thr Thr Leu Ile Met Leu Leu Ser Leu  
 60 65 70  
 Ala Ala Phe Ser Val Ile Ser Val Val Ser Tyr Leu Ile Leu Ala Leu  
 5 75 80 85 90  
 Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr Lys Ser Val Ile Gln Ala  
 95 100 105  
 Val Gln Lys Ser Glu Glu Gly His Pro Phe Lys Ala Tyr Leu Asp Val  
 110 115 120  
 10 Asp Ile Thr Leu Ser Ser Glu Ala Phe His Asn Tyr Met Asn Ala Ala  
 125 130 135  
 Met Val His Ile Asn Arg Ala Leu Lys Leu Ile Ile Arg Leu Phe Leu  
 140 145 150  
 Val Glu Asp Leu Val Asp Ser Leu Lys Leu Ala Val Phe Met Trp Leu  
 15 155 160 165 170  
 Met Thr Tyr Val Gly Ala Val Phe Asn Gly Ile Thr Leu Leu Ile Leu  
 175 180 185  
 Ala Glu Leu Leu Ile Phe Ser Val Pro Ile Val Tyr Glu Lys Tyr Lys  
 190 195 200  
 20 Thr Gln Ile Asp His Tyr Val Gly Ile Ala Arg Asp Gln Thr Lys Ser  
 205 210 215  
 Ile Val Glu Lys Ile Gln Ala Lys Leu Pro Gly Ile Ala Lys Lys Lys  
 220 225 230  
 Ala Glu  
 25 235  
  
 <210> 23  
 <211> 1069  
 <212> DNA  
 30 <213> Homo sapiens  
  
 <400> 23  
 agtgggaagac caggcagccc agctgaaggc agtaagctcg gctcacagtc gcaggagagt 60  
 tctgggggtac acggggcaaag gggcttgaga aggcccggag gcgaagccga agagaagcaa 120  
 35 ctgtgccccg gagaagagaa gctcgcccat tccagactgg gaaccagctt tcagtgaag 179

	atg gca ggg cca gaa ctg ttg ctc gac tcc aac atc cgc ctc tgg gtg	227
	Met Ala Gly Pro Glu Leu Leu Leu Asp Ser Asn Ile Arg Leu Trp Val	
	1                      5                      10                      15	
	gtc cta ccc atc gtt atc atc act ttc ttc gta ggc atg atc cgc cac	275
5	Val Leu Pro Ile Val Ile Ile Thr Phe Phe Val Gly Met Ile Arg His	
	20                      25                      30	
	tac gtg tcc atc ctg ctg cag agc gac aag aag ctc acc cag gaa caa	323
	Tyr Val Ser Ile Leu Leu Gln Ser Asp Lys Lys Leu Thr Gln Glu Gln	
	35                      40                      45	
10	gta tct gac agt caa gtc cta att cga agc aga gtc ctc agg gaa aat	371
	Val Ser Asp Ser Gln Val Leu Ile Arg Ser Arg Val Leu Arg Glu Asn	
	50                      55                      60	
	gga aaa tac att ccc aaa cag tct ttc ttg aca cga aaa tat tat ttc	419
	Gly Lys Tyr Ile Pro Lys Gln Ser Phe Leu Thr Arg Lys Tyr Tyr Phe	
15	65                      70                      75                      80	
	aac aac cca gag gat gga ttt ttc aaa aaa act aaa cgg aag gta gtg	467
	Asn Asn Pro Glu Asp Gly Phe Phe Lys Lys Thr Lys Arg Lys Val Val	
	85                      90                      95	
	cca cct tct cct atg act gat cct act atg ttg aca gac atg atg aaa	515
20	Pro Pro Ser Pro Met Thr Asp Pro Thr Met Leu Thr Asp Met Met Lys	
	100                      105                      110	
	ggg aat gta aca aat gtc ctc cct atg att ctt att ggt gga tgg atc	563
	Gly Asn Val Thr Asn Val Leu Pro Met Ile Leu Ile Gly Gly Trp Ile	
	115                      120                      125	
25	aac atg aca ttc tca ggc ttt gtc aca acc aag gtc cca ttt cca ctg	611
	Asn Met Thr Phe Ser Gly Phe Val Thr Thr Lys Val Pro Phe Pro Leu	
	130                      135                      140	
	acc ctc cgt ttt aag cct atg tta cag caa gga atc gag cta ctc aca	659
	Thr Leu Arg Phe Lys Pro Met Leu Gln Gln Gly Ile Glu Leu Leu Thr	
30	145                      150                      155                      160	
	tta gat gca tcc tgg gtg agt tct gca tcc tgg tac ttc ctc aat gta	707
	Leu Asp Ala Ser Trp Val Ser Ser Ala Ser Trp Tyr Phe Leu Asn Val	
	165                      170                      175	
	ttt ggg ctt cgg agc att tac tct ctg att ctg ggc caa gat aat gcc	755
35	Phe Gly Leu Arg Ser Ile Tyr Ser Leu Ile Leu Gly Gln Asp Asn Ala	

	180	185	190	
	gct gac caa tca cga atg atg cag gag cag atg acg gga gca gcc atg			803
	Ala Asp Gln Ser Arg Met Met Gln Glu Gln Met Thr Gly Ala Ala Met			
	195	200	205	
5	gcc atg ccc gca gac aca aac aaa gct ttc aag aca gag tgg gaa gct			851
	Ala Met Pro Ala Asp Thr Asn Lys Ala Phe Lys Thr Glu Trp Glu Ala			
	210	215	220	
	ttg gag ctg acg gat cac cag tgg gca cta gat gat gtc gaa gaa gag			899
	Leu Glu Leu Thr Asp His Gln Trp Ala Leu Asp Asp Val Glu Glu Glu			
10	225	230	235	240
	ctc atg gcc aaa gac ctc cac ttc gaa ggc atg ttc aaa aag gaa tta			947
	Leu Met Ala Lys Asp Leu His Phe Glu Gly Met Phe Lys Lys Glu Leu			
	245	250	255	
	cag acc tct att ttt tgaagaccga gcagggatta gctgtgtcag gaacttgg			1000
15	Gln Thr Ser Ile Phe			
	260			
	agttgcactt aaccttgtaa ctttgtttgg agctggcacc tottgaaata aaaaggagga			1060
	tgcacgagc			1069
20	<210> 24			
	<211> 261			
	<212> PRT			
	<213> Homo sapiens			
25	<400> 24			
	Met Ala Gly Pro Glu Leu Leu Leu Asp Ser Asn Ile Arg Leu Trp Val			
	1 5 10 15			
	Val Leu Pro Ile Val Ile Ile Thr Phe Phe Val Gly Met Ile Arg His			
	20 25 30			
30	Tyr Val Ser Ile Leu Leu Gln Ser Asp Lys Lys Leu Thr Gln Glu Gln			
	35 40 45			
	Val Ser Asp Ser Gln Val Leu Ile Arg Ser Arg Val Leu Arg Glu Asn			
	50 55 60			
	Gly Lys Tyr Ile Pro Lys Gln Ser Phe Leu Thr Arg Lys Tyr Tyr Phe			
35	65 70 75 80			

25

Asn Asn Pro Glu Asp Gly Phe Phe Lys Lys Thr Lys Arg Lys Val Val  
                     85                    90                    95  
 Pro Pro Ser Pro Met Thr Asp Pro Thr Met Leu Thr Asp Met Met Lys  
                     100                    105                    110  
 5 Gly Asn Val Thr Asn Val Leu Pro Met Ile Leu Ile Gly Gly Trp Ile  
                     115                    120                    125  
 Asn Met Thr Phe Ser Gly Phe Val Thr Thr Lys Val Pro Phe Pro Leu  
                     130                    135                    140  
 Thr Leu Arg Phe Lys Pro Met Leu Gln Gln Gly Ile Glu Leu Leu Thr  
 10 145                    150                    155                    160  
 Leu Asp Ala Ser Trp Val Ser Ser Ala Ser Trp Tyr Phe Leu Asn Val  
                     165                    170                    175  
 Phe Gly Leu Arg Ser Ile Tyr Ser Leu Ile Leu Gly Gln Asp Asn Ala  
                     180                    185                    190  
 15 Ala Asp Gln Ser Arg Met Met Gln Glu Gln Met Thr Gly Ala Ala Met  
                     195                    200                    205  
 Ala Met Pro Ala Asp Thr Asn Lys Ala Phe Lys Thr Glu Trp Glu Ala  
                     210                    215                    220  
 Leu Glu Leu Thr Asp His Gln Trp Ala Leu Asp Asp Val Glu Glu Glu  
 20 225                    230                    235                    240  
 Leu Met Ala Lys Asp Leu His Phe Glu Gly Met Phe Lys Lys Glu Leu  
                     245                    250                    255  
 Gln Thr Ser Ile Phe  
                     260

25

&lt;210&gt; 25

&lt;211&gt; 1759

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

30

&lt;400&gt; 25

attgtgcagc aggcggggccc ccgcgcggca gggccctgga cccgcgcggc tcccgggg  
 atg gtg agc aag gcg ctg ctg cgc ctc gtg tct gcc gtc aac cgc agg  
 Met Val Ser Lys Ala Leu Leu Arg Leu Val Ser Ala Val Asn Arg Arg

35

1

5

10

15

58

106

26.

	agg atg aag ctg ctg ctg ggc atc gcc ttg ctg gcc tac gtc gcc tct	154
	Arg Met Lys Leu Leu Leu Gly Ile Ala Leu Leu Ala Tyr Val Ala Ser	
	20 25 30	
	gtt tgg ggc aac ttc gtt aat atg agc ttt cta ctc aac agg tct atc	202
5	Val Trp Gly Asn Phe Val Asn Met Ser Phe Leu Leu Asn Arg Ser Ile	
	35 40 45	
	cag gaa aat ggt gaa cta aaa att gaa agc aag att gaa gag atg gtt	250
	Gln Glu Asn Gly Glu Leu Lys Ile Glu Ser Lys Ile Glu Glu Met Val	
	50 55 60	
10	gaa cca cta aga gag aaa atc aga gat tta gaa aaa agc ttt acc cag	298
	Glu Pro Leu Arg Glu Lys Ile Arg Asp Leu Glu Lys Ser Phe Thr Gln	
	65 70 75 80	
	aaa tac cca cca gta aag ttt tta tca gaa aag gat cgg aaa aga att	346
	Lys Tyr Pro Pro Val Lys Phe Leu Ser Glu Lys Asp Arg Lys Arg Ile	
15	85 90 95	
	ttg ata aca gga ggc gca ggg ttc gtg ggc tcc cat cta act gac aaa	394
	Leu Ile Thr Gly Gly Ala Gly Phe Val Gly Ser His Leu Thr Asp Lys	
	100 105 110	
	ctc atg atg gac ggc cac gag gtg acc gtg gtg gac aat ttc ttc acg	442
20	Leu Met Met Asp Gly His Glu Val Thr Val Val Asp Asn Phe Phe Thr	
	115 120 125	
	ggc agg aag aga aac gtg gag cac tgg atc gga cat gag aac ttc gag	490
	Gly Arg Lys Arg Asn Val Glu His Trp Ile Gly His Glu Asn Phe Glu	
	130 135 140	
25	ttg att aac cac gac gtg gtg gag ccc ctc tac atc gag ggc gtg gaa	538
	Leu Ile Asn His Asp Val Val Glu Pro Leu Tyr Ile Glu Gly Val Glu	
	145 150 155 160	
	gtg cga gtg gcc aga atc ttc aac acc ttt ggg cca cgc atg cac atg	586
	Val Arg Val Ala Arg Ile Phe Asn Thr Phe Gly Pro Arg Met His Met	
30	165 170 175	
	aac gat ggg cga gta gtc agc aac ttc atc ctg cag gcg ctc cag ggg	634
	Asn Asp Gly Arg Val Val Ser Asn Phe Ile Leu Gln Ala Leu Gln Gly	
	180 185 190	
	gag cca ctc acg gta tac gga tcc ggg tct cag aca agg gcg ttc cag	682
35	Glu Pro Leu Thr Val Tyr Gly Ser Gly Ser Gln Thr Arg Ala Phe Gln	

	195	200	205	
	tac gtc agc gat cta gtg aat ggc ctc gtg gct ctc atg aac agc aac			730
	Tyr Val Ser Asp Leu Val Asn Gly Leu Val Ala Leu Met Asn Ser Asn			
	210	215	220	
5	gtc agc agc ccg gtc aac ctg ggg aac cca gaa gaa cac aca atc cta			778
	Val Ser Ser Pro Val Asn Leu Gly Asn Pro Glu Glu His Thr Ile Leu			
	225	230	235	240
	gaa ttt gct cag tta att aaa aac ctt gtt ggt agc gga agt gaa att			826
	Glu Phe Ala Gln Leu Ile Lys Asn Leu Val Gly Ser Gly Ser Glu Ile			
10	245	250	255	
	cag ttt ctc tcc gaa gcc cag gat gac cca cag aaa aga aaa cca gac			874
	Gln Phe Leu Ser Glu Ala Gln Asp Asp Pro Gln Lys Arg Lys Pro Asp			
	260	265	270	
	atc aaa aaa gca aag ctg atg ctg ggg tgg gag ccc gtg gtc ccg ctg			922
15	Ile Lys Lys Ala Lys Leu Met Leu Gly Trp Glu Pro Val Val Pro Leu			
	275	280	285	
	gag gaa ggt tta aac aaa gca att cac tac ttc cgt aaa gaa ctc gag			970
	Glu Glu Gly Leu Asn Lys Ala Ile His Tyr Phe Arg Lys Glu Leu Glu			
	290	295	300	
20	tac cag gca aat aat cag tac atc ccc aaa cca aag cct gcc aga ata			1018
	Tyr Gln Ala Asn Asn Gln Tyr Ile Pro Lys Pro Lys Pro Ala Arg Ile			
	305	310	315	320
	aag aaa gga cgg act cgc cac agc tgaactcctc acttttagga cacaagac			1070
	Lys Lys Gly Arg Thr Arg His Ser			
25	325			
	taccattgta cacttgatgg gatgtatttt tggttttttt ttgttgctgt ttaaagaaag			1130
	actttaacag gtgtcatgaa gaacaaactg gaatttcatt ctgaagcttg ctttaatgaa			1190
	atggatgtgc ctaaaagctc ccctcaaaaa actgcagatt ttgccttgca ctttttgaat			1250
	ctctcttttt atgtaaaata gcgtagatgc atctctgcgt attttcaagt ttttttatct			1310
30	tgctgtgaga gcatatgttg tgactgtcgt tgacagtttt atttactggt ttctttgtga			1370
	agctgaaaag gaacattaag cgggacaaaa aatgccgatt ttatttataa aagtgggtac			1430
	ttaataaatg agtcgttata ctatgcataa agaaaaatcc tagcagtatt gtcaggtggt			1490
	ggtgcgccgg cattgatttt agggcagata aaagaattct gtgtgagagc tttatgtttc			1550
	tcttttaatt cagagttttt ccaaggtcta cttttgagtt gcaaacttga ctttgaaata			1610
35	ttcctgttgg tcatgatcaa ggatatttga aatcactact gtgttttgct gcgtatctgg			1670



ggcgggggca ggttgggggg cacaaagtta acatattctt ggttaaccat ggttaaatat 1730  
gctatttttaa taaaatattg aaactcacc 1759

<210> 26

5 <211> 328

<212> PRT

<213> Homo sapiens

<400> 26

10 Met Val Ser Lys Ala Leu Leu Arg Leu Val Ser Ala Val Asn Arg Arg  
1 5 10 15  
Arg Met Lys Leu Leu Leu Gly Ile Ala Leu Leu Ala Tyr Val Ala Ser  
20 25 30  
Val Trp Gly Asn Phe Val Asn Met Ser Phe Leu Leu Asn Arg Ser Ile  
15 35 40 45  
Gln Glu Asn Gly Glu Leu Lys Ile Glu Ser Lys Ile Glu Glu Met Val  
50 55 60  
Glu Pro Leu Arg Glu Lys Ile Arg Asp Leu Glu Lys Ser Phe Thr Gln  
65 70 75 80  
20 Lys Tyr Pro Pro Val Lys Phe Leu Ser Glu Lys Asp Arg Lys Arg Ile  
85 90 95  
Leu Ile Thr Gly Gly Ala Gly Phe Val Gly Ser His Leu Thr Asp Lys  
100 105 110  
Leu Met Met Asp Gly His Glu Val Thr Val Val Asp Asn Phe Phe Thr  
25 115 120 125  
Gly Arg Lys Arg Asn Val Glu His Trp Ile Gly His Glu Asn Phe Glu  
130 135 140  
Leu Ile Asn His Asp Val Val Glu Pro Leu Tyr Ile Glu Gly Val Glu  
145 150 155 160  
30 Val Arg Val Ala Arg Ile Phe Asn Thr Phe Gly Pro Arg Met His Met  
165 170 175  
Asn Asp Gly Arg Val Val Ser Asn Phe Ile Leu Gln Ala Leu Gln Gly  
180 185 190  
Glu Pro Leu Thr Val Tyr Gly Ser Gly Ser Gln Thr Arg Ala Phe Gln  
35 195 200 205

Tyr Val Ser Asp Leu Val Asn Gly Leu Val Ala Leu Met Asn Ser Asn  
 210 215 220  
 Val Ser Ser Pro Val Asn Leu Gly Asn Pro Glu Glu His Thr Ile Leu  
 225 230 235 240  
 5 Glu Phe Ala Gln Leu Ile Lys Asn Leu Val Gly Ser Gly Ser Glu Ile  
 245 250 255  
 Gln Phe Leu Ser Glu Ala Gln Asp Asp Pro Gln Lys Arg Lys Pro Asp  
 260 265 270  
 Ile Lys Lys Ala Lys Leu Met Leu Gly Trp Glu Pro Val Val Pro Leu  
 10 275 280 285  
 Glu Glu Gly Leu Asn Lys Ala Ile His Tyr Phe Arg Lys Glu Leu Glu  
 290 295 300  
 Tyr Gln Ala Asn Asn Gln Tyr Ile Pro Lys Pro Lys Pro Ala Arg Ile  
 305 310 315 320  
 15 Lys Lys Gly Arg Thr Arg His Ser  
 325

<210> 27  
 <211> 1697  
 20 <212> DNA  
 <213> Homo sapiens

<400> 27  
 aaaaggatac gggagttcct ccttgctctc gccctactc tttctggtgt tagatcgagc 60  
 25 taccctctaa aagcagttta gagtggtaaa aaaaaaaaaa aacacaccaa acgctcgag 120  
 ccacaaaagg g atg aaa ttt ctt ctg gac atc ctc ctg ctt ctc ccg tta 170  
 Met Lys Phe Leu Leu Asp Ile Leu Leu Leu Leu Pro Leu  
 1 5 10  
 ctg atc gtc tgc tcc cta gag tcc ttc gtg aag ctt ttt att cct aag 218  
 30 Leu Ile Val Cys Ser Leu Glu Ser Phe Val Lys Leu Phe Ile Pro Lys  
 15 20 25  
 agg aga aaa tca gtc acc ggc gaa atc gtg ctg att aca gga gct ggg 266  
 Arg Arg Lys Ser Val Thr Gly Glu Ile Val Leu Ile Thr Gly Ala Gly  
 30 35 40 45  
 35 cat gga att ggg aga ctg act gcc tat gaa ttt gct aaa ctt aaa agc 314

30.

	His Gly Ile Gly Arg Leu Thr Ala Tyr Glu Phe Ala Lys Leu Lys Ser	
	50 55 60	
	aag ctg gtt ctc tgg gat ata aat aag cat gga ctg gag gaa aca gct	362
	Lys Leu Val Leu Trp Asp Ile Asn Lys His Gly Leu Glu Glu Thr Ala	
5	65 70 75	
	gcc aaa tgc aag gga ctg ggt gcc aag gtt cat acc ttt gtg gta gac	410
	Ala Lys Cys Lys Gly Leu Gly Ala Lys Val His Thr Phe Val Val Asp	
	80 85 90	
	tgc agc aac cga gaa gat att tac agc tct gca aag aag gtg aag gca	458
10	Cys Ser Asn Arg Glu Asp Ile Tyr Ser Ser Ala Lys Lys Val Lys Ala	
	95 100 105	
	gaa att gga gat gtt agt att tta gta aat aat gct ggt gta gtc tat	506
	Glu Ile Gly Asp Val Ser Ile Leu Val Asn Asn Ala Gly Val Val Tyr	
	110 115 120 125	
15	aca tca gat ttg ttt gct aca caa gat cct cag att gaa aag act ttt	554
	Thr Ser Asp Leu Phe Ala Thr Gln Asp Pro Gln Ile Glu Lys Thr Phe	
	130 135 140	
	gaa gtt aat gta ctt gca cat ttc tgg act aca aag gca ttt ctt cct	602
	Glu Val Asn Val Leu Ala His Phe Trp Thr Thr Lys Ala Phe Leu Pro	
20	145 150 155	
	gca atg acg aag aat aac cat ggc cat att gtc act gtg gct tcg gca	650
	Ala Met Thr Lys Asn Asn His Gly His Ile Val Thr Val Ala Ser Ala	
	160 165 170	
	gct gga cat gtc tcg gtc ccc ttc tta ctg gct tac tgt tca agc aag	698
25	Ala Gly His Val Ser Val Pro Phe Leu Leu Ala Tyr Cys Ser Ser Lys	
	175 180 185	
	ttt gct gct gtt gga ttt cat aaa act ttg aca gat gaa ctg gct gcc	746
	Phe Ala Ala Val Gly Phe His Lys Thr Leu Thr Asp Glu Leu Ala Ala	
	190 195 200 205	
30	tta caa ata act gga gtc aaa aca aca tgt ctg tgt cct aat ttc gta	794
	Leu Gln Ile Thr Gly Val Lys Thr Thr Cys Leu Cys Pro Asn Phe Val	
	210 215 220	
	aac act ggc ttc atc aaa aat cca agt aca agt ttg gga ccc act ctg	842
	Asn Thr Gly Phe Ile Lys Asn Pro Ser Thr Ser Leu Gly Pro Thr Leu	
35	225 230 235	

	gaa cct gag gaa gtg gta aac agg ctg atg cat ggg att ctg act gag	890
	Glu Pro Glu Glu Val Val Asn Arg Leu Met His Gly Ile Leu Thr Glu	
	240 245 250	
	cag aag atg att ttt att cca tct tct ata gct ttt tta aca aca ttg	938
5	Gln Lys Met Ile Phe Ile Pro Ser Ser Ile Ala Phe Leu Thr Thr Leu	
	255 260 265	
	gaa agg atc ctt cct gag cgt ttc ctg gca gtt tta aaa cga aaa atc	986
	Glu Arg Ile Leu Pro Glu Arg Phe Leu Ala Val Leu Lys Arg Lys Ile	
	270 275 280 285	
10	agt gtt aag ttt gat gca gtt att gga tat aaa atg aaa gcg caa	1031
	Ser Val Lys Phe Asp Ala Val Ile Gly Tyr Lys Met Lys Ala Gln	
	290 295 300	
	taagcacct agttttctga aaactgattt accagggttta ggttgatgtc atctaatagt	1090
	gccagaattt taatgtttga acttctgttt tttctaatta tccccatttc ttcaatatca	1150
15	tttttgaggc tttggcagtc ttcatttact accacttggt ctttagccaa aagctgatta	1210
	catatgatat aaacagagaa ataccttttag aggtgacttt aaggaaaatg aagaaaaaga	1270
	accaaaatga ctttattaaa ataatttcca agattatttg tggctcacct gaaggctttg	1330
	caaaatttgt accataaccg tttattttaac atatattttt atttttgatt gcacttaa	1390
	tttgataat ttgtgtttct ttttctgttc tacataaaat cagaaacttc aagctctcta	1450
20	aataaaatga aggactatat ctagtggtat ttcacaatga atatcatgaa ctctcaatgg	1510
	gtaggtttca tcctacccat tgccactctg tttcctgaga gatacctcac attccaatgc	1570
	caaacatttc tgcacagga agctagaggt ggatacacgt gttgcaagta taaaagcatc	1630
	actgggattt aaggagaatt gagagaatgt accacaaaat ggcagcaata ataaatggat	1690
	cacactt	1697
25	<210> 28	
	<211> 300	
	<212> PRT	
	<213> Homo sapiens	
30	<400> 28	
	Met Lys Phe Leu Leu Asp Ile Leu Leu Leu Leu Pro Leu	
	1 5 10	
	Leu Ile Val Cys Ser Leu Glu Ser Phe Val Lys Leu Phe Ile Pro Lys	
35	15 20 25	

Arg Arg Lys Ser Val Thr Gly Glu Ile Val Leu Ile Thr Gly Ala Gly  
 30 35 40 45  
 His Gly Ile Gly Arg Leu Thr Ala Tyr Glu Phe Ala Lys Leu Lys Ser  
 50 55 60  
 5 Lys Leu Val Leu Trp Asp Ile Asn Lys His Gly Leu Glu Glu Thr Ala  
 65 70 75  
 Ala Lys Cys Lys Gly Leu Gly Ala Lys Val His Thr Phe Val Val Asp  
 80 85 90  
 Cys Ser Asn Arg Glu Asp Ile Tyr Ser Ser Ala Lys Lys Val Lys Ala  
 10 95 100 105  
 Glu Ile Gly Asp Val Ser Ile Leu Val Asn Asn Ala Gly Val Val Tyr  
 110 115 120 125  
 Thr Ser Asp Leu Phe Ala Thr Gln Asp Pro Gln Ile Glu Lys Thr Phe  
 130 135 140  
 15 Glu Val Asn Val Leu Ala His Phe Trp Thr Thr Lys Ala Phe Leu Pro  
 145 150 155  
 Ala Met Thr Lys Asn Asn His Gly His Ile Val Thr Val Ala Ser Ala  
 160 165 170  
 Ala Gly His Val Ser Val Pro Phe Leu Leu Ala Tyr Cys Ser Ser Lys  
 20 175 180 185  
 Phe Ala Ala Val Gly Phe His Lys Thr Leu Thr Asp Glu Leu Ala Ala  
 190 195 200 205  
 Leu Gln Ile Thr Gly Val Lys Thr Thr Cys Leu Cys Pro Asn Phe Val  
 210 215 220  
 25 Asn Thr Gly Phe Ile Lys Asn Pro Ser Thr Ser Leu Gly Pro Thr Leu  
 225 230 235  
 Glu Pro Glu Glu Val Val Asn Arg Leu Met His Gly Ile Leu Thr Glu  
 240 245 250  
 Gln Lys Met Ile Phe Ile Pro Ser Ser Ile Ala Phe Leu Thr Thr Leu  
 30 255 260 265  
 Glu Arg Ile Leu Pro Glu Arg Phe Leu Ala Val Leu Lys Arg Lys Ile  
 270 275 280 285  
 Ser Val Lys Phe Asp Ala Val Ile Gly Tyr Lys Met Lys Ala Gln  
 290 295 300

&lt;210&gt; 29

&lt;211&gt; 814

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

5

&lt;400&gt; 29

```

agaatccccgg acagccctgc tccctgcagc caggtgtagt ttcgggagcc actggggcca      60
aagtgagagt ccagcgggtct tccagcgctt gggccacggc ggcggccctg ggagcagagg      120
tggagcgacc ccattacgct aaag atg aaa ggc tgg ggt tgg ctg gcc ctg      171
                                Met Lys Gly Trp Gly Trp Leu Ala Leu
                                1               5

ctt ctg ggg gcc ctg ctg gga acc gcc tgg gct cgg agg agc cag gat      219
Leu Leu Gly Ala Leu Leu Gly Thr Ala Trp Ala Arg Arg Ser Gln Asp
    10               15               20               25

ctc cac tgt gga gca tgc agg gct ctg gtg gat gaa cta gaa tgg gaa      267
Leu His Cys Gly Ala Cys Arg Ala Leu Val Asp Glu Leu Glu Trp Glu
                30               35               40

att gcc cag gtg gac ccc aag aag acc att cag atg gga tct ttc cgg      315
Ile Ala Gln Val Asp Pro Lys Lys Thr Ile Gln Met Gly Ser Phe Arg
                45               50               55

atc aat cca gat ggc agc cag tca gtg gtg gag gtg cct tat gcc cgc      363
Ile Asn Pro Asp Gly Ser Gln Ser Val Val Glu Val Pro Tyr Ala Arg
                60               65               70

tca gag gcc cac ctc aca gag ctg ctg gag gag ata tgt gac cgg atg      411
Ser Glu Ala His Leu Thr Glu Leu Leu Glu Glu Ile Cys Asp Arg Met
    75               80               85

aag gag tat ggg gaa cag att gat cct tcc acc cat cgc aag aac tac      459
Lys Glu Tyr Gly Glu Gln Ile Asp Pro Ser Thr His Arg Lys Asn Tyr
    90               95               100               105

gta cgt gta gtg ggc cgg aat gga gaa tcc agt gaa ctg gac cta caa      507
Val Arg Val Val Gly Arg Asn Gly Glu Ser Ser Glu Leu Asp Leu Gln
                110               115               120

ggc atc cga atc gac tca gat att agc ggc acc ctc aag ttt gcg tgt      555
Gly Ile Arg Ile Asp Ser Asp Ile Ser Gly Thr Leu Lys Phe Ala Cys
    125               130               135

```

gag agc att gtg gag gaa tac gag gat gaa ctc att gaa ttc ttt tcc 603  
 Glu Ser Ile Val Glu Glu Tyr Glu Asp Glu Leu Ile Glu Phe Phe Ser  
 140 145 150  
 cga gag gct gac aat gtt aaa gac aaa ctt tgc agt aag cga aca gat 651  
 5 Arg Glu Ala Asp Asn Val Lys Asp Lys Leu Cys Ser Lys Arg Thr Asp  
 155 160 165  
 ctt tgt gac cat gcc ctg cac ata tcg cat gat gag cta tgaaccactg 700  
 Leu Cys Asp His Ala Leu His Ile Ser His Asp Glu Leu  
 170 175 180  
 10 gagcagccca cactggcttg atggatcacc cccaggaggg gaaaatggtg gcaatgcctt 760  
 ttatatatta tgtttttact gaaattaact gaaaaaatat gaaacccaaaa gtac 814  
  
 <210> 30  
 <211> 182  
 15 <212> PRT  
 <213> Homo sapiens  
  
 <400> 30  
  
 Met Lys Gly Trp Gly Trp Leu Ala Leu  
 20 1 5  
 Leu Leu Gly Ala Leu Leu Gly Thr Ala Trp Ala Arg Arg Ser Gln Asp  
 10 15 20 25  
 Leu His Cys Gly Ala Cys Arg Ala Leu Val Asp Glu Leu Glu Trp Glu  
 30 35 40  
 25 Ile Ala Gln Val Asp Pro Lys Lys Thr Ile Gln Met Gly Ser Phe Arg  
 45 50 55  
 Ile Asn Pro Asp Gly Ser Gln Ser Val Val Glu Val Pro Tyr Ala Arg  
 60 65 70  
 Ser Glu Ala His Leu Thr Glu Leu Leu Glu Glu Ile Cys Asp Arg Met  
 30 75 80 85  
 Lys Glu Tyr Gly Glu Gln Ile Asp Pro Ser Thr His Arg Lys Asn Tyr  
 90 95 100 105  
 Val Arg Val Val Gly Arg Asn Gly Glu Ser Ser Glu Leu Asp Leu Gln  
 110 115 120  
 35 Gly Ile Arg Ile Asp Ser Asp Ile Ser Gly Thr Leu Lys Phe Ala Cys

35.

	125	130	135	
	Glu Ser Ile Val Glu Glu Tyr Glu Asp Glu Leu Ile Glu Phe Phe Ser			
	140	145	150	
	Arg Glu Ala Asp Asn Val Lys Asp Lys Leu Cys Ser Lys Arg Thr Asp			
5	155	160	165	
	Leu Cys Asp His Ala Leu His Ile Ser His Asp Glu Leu			
	170	175	180	
	<210> 31			
10	<211> 511			
	<212> DNA			
	<213> Homo sapiens			
	<400> 31			
15	gttacgaagc tgcaggagcg ag atg gag gtg gac gca ccg ggt gtt gat ggt			52
	Met Glu Val Asp Ala Pro Gly Val Asp Gly			
	1	5	10	
	CGA GAT GGT CTC CGG GAG CGG CGA GGC TTT AGC GAG GGA GGG AGG CAG			100
	Arg Asp Gly Leu Arg Glu Arg Arg Gly Phe Ser Glu Gly Gly Arg Gln			
20	15	20	25	
	aac ttc gat gtg agg cct cag tct ggg gca aat ggg ctt ccc aaa cac			148
	Asn Phe Asp Val Arg Pro Gln Ser Gly Ala Asn Gly Leu Pro Lys His			
	30	35	40	
	tcc tac tgg ttg gac ctc tgg ctt ttc atc ctt ttc gat gtg gtg gtg			196
25	Ser Tyr Trp Leu Asp Leu Trp Leu Phe Ile Leu Phe Asp Val Val Val			
	45	50	55	
	ttt ctc ttt gtg tat ttt ttg cca tgacttggtc gctgatatct aaattaagaa			250
	Phe Leu Phe Val Tyr Phe Leu Pro			
	60	65		
30	gttggttctt gactgaattc tgaaaatggc tacaaacttc ttgaataaag aagacaggac			310
	tctcaataga agaatttcac atctccaagg gacccttcct ttcattttac actttgttac			370
	taatttgcag aactctatta attgggtagg atttcaccca ttcctagcta agttcttaaa			430
	attaaaccct ttggttcgtg tttaaaaact ttcaaacatc tgatggcttt acaggggctg			490
	aatataaaag catttgtact t			511
35				



<210> 32

<211> 66

<212> PRT

<213> Homo sapiens

5

<400> 32

[illegible]

<210> 33

**<211> 1126**

20           <212> DNA

<213> Homo sapiens

<400> 33

	ctcttcacgg agccgcgcgg ctgcgggggc gcaaataaggg tcaactgggcc gcttggcggt	60
25	gtcgttgcgg taccaggtcc gcgtgagggg ttcggggggtt ctgggcaggc aca atg	116
	Met	
	1	
	gcg tct cga gca ggc ccg cga gcg gcc ggc acc gac ggc agc gac ttt	164
	Ala Ser Arg Ala Gly Pro Arg Ala Ala Gly Thr Asp Gly Ser Asp Phe	
30	5 10 15	
	cag cac cgg gag cgc gtc gcc atg cac tac cag atg agt gtg acc ctc	212
	Gln His Arg Glu Arg Val Ala Met His Tyr Gln Met Ser Val Thr Leu	
	20 25 30	
	aag tat gaa atc aag aag ctg atc tac gta cat ctg gtc ata tgg ctg	260
35	Lys Tyr Glu Ile Lys Lys Leu Ile Tyr Val His Leu Val Ile Trp Leu	

	35	40	45	
	ctg ctg gtt gct aag atg agc gtg gga cac ctg agg ctc ttg tca cat			308
	Leu Leu Val Ala Lys Met Ser Val Gly His Leu Arg Leu Leu Ser His			
	50	55	60	65
5	gat cag gtg gcc atg ccc tat cag tgg gaa tac ccg tat ttg ctg agc			356
	Asp Gln Val Ala Met Pro Tyr Gln Trp Glu Tyr Pro Tyr Leu Leu Ser			
	70	75	80	
	att ttg ccc tct ctc ttg ggc ctt ctc tcc ttt ccc cgc aac aac att			404
	Ile Leu Pro Ser Leu Leu Gly Leu Leu Ser Phe Pro Arg Asn Asn Ile			
10	85	90	95	
	agc tac ctg gtg ctc tcc atg atc agc atg gga ctc ttt tcc atc gct			452
	Ser Tyr Leu Val Leu Ser Met Ile Ser Met Gly Leu Phe Ser Ile Ala			
	100	105	110	
	cca ctc att tat ggc agc atg gag atg ttc cct gct gca cag cag ctc			500
15	Pro Leu Ile Tyr Gly Ser Met Glu Met Phe Pro Ala Ala Gln Gln Leu			
	115	120	125	
	tac cgc cat ggc aag gcc tac cgt ttc ctc ttt ggt ttt tct gcc gtt			548
	Tyr Arg His Gly Lys Ala Tyr Arg Phe Leu Phe Gly Phe Ser Ala Val			
	130	135	140	145
20	tcc atc atg tac ctg gtg ttg gtg ttg gca gtg caa gtg cat gcc tgg			596
	Ser Ile Met Tyr Leu Val Leu Val Leu Ala Val Gln Val His Ala Trp			
	150	155	160	
	cag ttg tac tac agc aag aag ctc cta gac tct tgg ttc acc agc aca			644
	Gln Leu Tyr Tyr Ser Lys Lys Leu Leu Asp Ser Trp Phe Thr Ser Thr			
25	165	170	175	
	cag gag aag aag cat aaa tgaagcctct ttgggggtgaa gcctggacat cccatcga			700
	Gln Glu Lys Lys His Lys			
	180			
	atgaaaggac actagtacag cggttccaaa atcccttctg gtgattttag cagctgtgat			760
30	gttggtacct ggtgcagacc aggccaaagt tctggaaagc tccttttgcc atctgctgag			820
	gtggcaaaac tataatttat tcctggttgg ctagaactgg gtgaccgaca gctatgaaac			880
	aaatttcagc tgtttgaagt tgaactttga ggtttttctt taagaatgag cttcgtcctt			940
	gcctctactc ggtcattctc cccatttcca tccattaccc cttagccatt gagactaaag			1000
	gaaatagggg ataaatcaaa ttacttcatc tctaggtcac gggtcaggaa acatttgggc			1060
35	agctgctccc ttggcagctg tgggtctcctc tgcaaagcat ttaattaaa aacctcaata			1120

aagatg

1126

&lt;210&gt; 34

&lt;211&gt; 183

5 &lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

Met

10

1

Ala Ser Arg Ala Gly Pro Arg Ala Ala Gly Thr Asp Gly Ser Asp Phe

5

10

15

Gln His Arg Glu Arg Val Ala Met His Tyr Gln Met Ser Val Thr Leu

20

25

30

15

Lys Tyr Glu Ile Lys Lys Leu Ile Tyr Val His Leu Val Ile Trp Leu

35

40

45

Leu Leu Val Ala Lys Met Ser Val Gly His Leu Arg Leu Leu Ser His

50

55

60

65

Asp Gln Val Ala Met Pro Tyr Gln Trp Glu Tyr Pro Tyr Leu Leu Ser

20

70

75

80

Ile Leu Pro Ser Leu Leu Gly Leu Leu Ser Phe Pro Arg Asn Asn Ile

85

90

95

Ser Tyr Leu Val Leu Ser Met Ile Ser Met Gly Leu Phe Ser Ile Ala

100

105

110

25

Pro Leu Ile Tyr Gly Ser Met Glu Met Phe Pro Ala Ala Gln Gln Leu

115

120

125

Tyr Arg His Gly Lys Ala Tyr Arg Phe Leu Phe Gly Phe Ser Ala Val

130

135

140

145

Ser Ile Met Tyr Leu Val Leu Val Leu Ala Val Gln Val His Ala Trp

30

150

155

160

Gln Leu Tyr Tyr Ser Lys Lys Leu Leu Asp Ser Trp Phe Thr Ser Thr

165

170

175

Gln Glu Lys Lys His Lys

180

35

&lt;210&gt; 35

&lt;211&gt; 2015

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

5

&lt;400&gt; 35

	atgtagtgag accctcgcca ggtctgagag tcaactggagc taccagaagc atc atg	56
	Met	
	1	
10	ggg ccc tgg gga gag cca gag ctc ctg gtg tgg cgc ccc gag gcg gta	104
	Gly Pro Trp Gly Glu Pro Glu Leu Leu Val Trp Arg Pro Glu Ala Val	
	5 10 15	
	gct tca gag cct cca gtg cct gtg ggg ctg gag gtg aag ttg ggg gcc	152
	Ala Ser Glu Pro Pro Val Pro Val Gly Leu Glu Val Lys Leu Gly Ala	
15	20 25 30	
	ctg gtg ctg ctg ctg gtg ctc acc ctc ctc tgc agc ctg gtg ccc atc	200
	Leu Val Leu Leu Leu Val Leu Thr Leu Leu Cys Ser Leu Val Pro Ile	
	35 40 45	
	tgt gtg ctg cgc cgg cca gga gct aac cat gaa ggc tca gct tcc cgc	248
20	Cys Val Leu Arg Arg Pro Gly Ala Asn His Glu Gly Ser Ala Ser Arg	
	50 55 60 65	
	cag aaa gcc ctg agc cta gta agc tgt ttc gcg ggg ggc gtc ttt ttg	296
	Gln Lys Ala Leu Ser Leu Val Ser Cys Phe Ala Gly Gly Val Phe Leu	
	70 75 80	
25	gcc act tgt ctc ctg gac ctg ctg cct gac tac ctg gct gcc ata gat	344
	Ala Thr Cys Leu Leu Asp Leu Leu Pro Asp Tyr Leu Ala Ala Ile Asp	
	85 90 95	
	gag gcc ctg gca gcc ttg cac gtg acg ctc cag ttc cca ctg caa gag	392
	Glu Ala Leu Ala Ala Leu His Val Thr Leu Gln Phe Pro Leu Gln Glu	
30	100 105 110	
	ttc atc ctg gcc atg ggc ttc ttc ctg gtc ctg gtg atg gag cag atc	440
	Phe Ile Leu Ala Met Gly Phe Phe Leu Val Leu Val Met Glu Gln Ile	
	115 120 125	
	aca ctg gct tac aag gag cag tca ggg ccg tca cct ctg gag gaa aca	488
35	Thr Leu Ala Tyr Lys Glu Gln Ser Gly Pro Ser Pro Leu Glu Glu Thr	

	130	135	140	145	
	agg gct ctg ctg gga aca gtg aat ggt ggg ccg cag cat tgg cat gat	536			
	Arg Ala Leu Leu Gly Thr Val Asn Gly Gly Pro Gln His Trp His Asp				
	150	155	160		
5	ggg cca ggg gtc cca cag gcg agt gga gcc cca gca acc ccc tca gcc	584			
	Gly Pro Gly Val Pro Gln Ala Ser Gly Ala Pro Ala Thr Pro Ser Ala				
	165	170	175		
	ttg cgt gcc tgt gta ctg gtg ttc tcc ctg gcc ctc cac tcc gtg ttc	632			
	Leu Arg Ala Cys Val Leu Val Phe Ser Leu Ala Leu His Ser Val Phe				
10	180	185	190		
	gag ggg ctg gcg gta ggg ctg cag cga gac cgg gct cgg gcc atg gag	680			
	Glu Gly Leu Ala Val Gly Leu Gln Arg Asp Arg Ala Arg Ala Met Glu				
	195	200	205		
	ctg tgc ctg gct ttg ctg ctc cac aag ggc atc ctg gct gtc agc ctg	728			
15	Leu Cys Leu Ala Leu Leu Leu His Lys Gly Ile Leu Ala Val Ser Leu				
	210	215	220	225	
	tcc ctg cgg ctg ttg cag agc cac ctt agg gca cag gtg gtg gct ggc	776			
	Ser Leu Arg Leu Leu Gln Ser His Leu Arg Ala Gln Val Val Ala Gly				
	230	235	240		
20	tgt ggg atc ctc ttc tca tgc atg aca cct cta ggc atc ggg ctg ggt	824			
	Cys Gly Ile Leu Phe Ser Cys Met Thr Pro Leu Gly Ile Gly Leu Gly				
	245	250	255		
	gca gct ctg gca gag tcg gca gga cct ctg cac cag ctg gcc cag tct	872			
	Ala Ala Leu Ala Glu Ser Ala Gly Pro Leu His Gln Leu Ala Gln Ser				
25	260	265	270		
	gtg cta gag ggc atg gca gct ggc acc ttt ctc tat atc acc ttt ctg	920			
	Val Leu Glu Gly Met Ala Ala Gly Thr Phe Leu Tyr Ile Thr Phe Leu				
	275	280	285		
	gaa atc ctg ccc cag gag ctg gcc agt tct gag caa agg atc ctc aag	968			
30	Glu Ile Leu Pro Gln Glu Leu Ala Ser Ser Glu Gln Arg Ile Leu Lys				
	290	295	300	305	
	gtc att ctg ctc cta gca ggc ttt gcc ctg ctc act ggc ctg ctc ttc	1016			
	Val Ile Leu Leu Leu Ala Gly Phe Ala Leu Leu Thr Gly Leu Leu Phe				
	310	315	320		
35	atc caa atc tagggggctt caagagaggg gcaggggaga ttgatgatca ggtgc	1070			

Ile Gln Ile

5 ccctgttctc ccttcctcc cccagttgtg gggaatagga aggaaagggg aagggaata 1130  
 ctgaggacca aaaagttctc tgggagctaa agatagagcc tttggggcta tctgactaat 1190  
 gagaggggaag tgggcagaca agaggctggc cccagtccca aggaacaaga gatggtcaag 1250  
 tcgctagaga catatcaggg gacattagga ttggggaaga cacttgactg ctagaatcag 1310  
 aggttggaaca ctatacataa ggacaggctc acatgggagg ctggaggtgg gtaccagct 1370  
 gctgtggaac gggatatggac aggtcataaa cctagagtca gtgtcctgtt ggtcctagcc 1430  
 catttcagca ccctgccact tggagtggac ccctcctact cttcttagcg cctaccctca 1490  
 10 tacctatctc cctcctccca tctcctaggg gactggcgcc aaatgggtctc tcctgccaa 1550  
 ttttggatc ttctctggcc tctccagtc tgcttactcc tctattttta aagtgccaaa 1610  
 caaatcccct tcctctttct caaagcacag taatgtggca ctgagcccta cccagcacct 1670  
 cagtgaaggg ggctgcttg ctctttattt tggccccgga tcctgggggtg gggcagaaat 1730  
 attttctggg ctggggtagg aggaagggtg ttgcagccat ctactgctgc tgtaccctag 1790  
 15 gaatatgggg acatggacat ggtgtcccat gcccagatga taaacactga gctgccccaa 1850  
 cattttttta aatacacccg aggagcccaa gggggaaggg caatgcctac cccagcggt 1910  
 atttttgggg agggagggt gtgcataggg acatattctt tagaatctat tttattaact 1970  
 gacctgtttt gggacctgtt acccaaataa aagatgtttc tagac 2015

20 <210> 36  
 <211> 324  
 <212> PRT  
 <213> Homo sapiens

25 <400> 36

Met

1

Gly Pro Trp Gly Glu Pro Glu Leu Leu Val Trp Arg Pro Glu Ala Val  
 5 10 15  
 30 Ala Ser Glu Pro Pro Val Pro Val Gly Leu Glu Val Lys Leu Gly Ala  
 20 25 30  
 Leu Val Leu Leu Leu Val Leu Thr Leu Leu Cys Ser Leu Val Pro Ile  
 35 40 45  
 Cys Val Leu Arg Arg Pro Gly Ala Asn His Glu Gly Ser Ala Ser Arg  
 35 50 55 60 65



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>6</sup> :</b> <b>C12N 15/12, C07K 14/47, C12N 5/10</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/55862</b>
		<b>(43) International Publication Date:</b> 4 November 1999 (04.11.99)
<b>(21) International Application Number:</b> PCT/JP99/02226 <b>(22) International Filing Date:</b> 27 April 1999 (27.04.99)  <b>(30) Priority Data:</b> 10/119395      28 April 1998 (28.04.98)      JP  <b>(71) Applicants (for all designated States except US):</b> SAGAMI CHEMICAL RESEARCH CENTER [JP/JP]; 4-1, Nishi-Ohnuma 4-chome, Sagamihara-shi, Kanagawa 229-0012 (JP). PROTEGENE INC. [JP/JP]; 2-20-3, Naka-cho, Meguro-ku, Tokyo 153-0065 (JP).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> <u>KATO</u> , Seishi [JP/JP]; 3-46-50, Wakamatsu, Sagamihara-shi, Kanagawa 229-0014 (JP). <u>KIMURA</u> , Tomoko [JP/JP]; 302, 4-1-28, Nishiikuta, Tama-ku, Kawasaki-shi, Kanagawa 214-0037 (JP).  <b>(74) Agents:</b> AOYAMA, Tamotsu et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).		<b>(81) Designated States:</b> AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 2 March 2000 (02.03.00)
<b>(54) Title:</b> HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAs ENCODING THESE PROTEINS		
<b>(57) Abstract</b>  A protein comprising any of the amino acid sequences represented by Sequence Nos. 1 to 9, a DNA coding for said protein, exemplified by a cDNA comprising any of the base sequences represented by Sequence Nos. 10 to 18, and an expression vector of said cDNA as well as an eucaryotic cell expressing said cDNA. Said protein and eucaryotic cell having said protein on the membrane surface can be provided by expression of a cDNA coding for a human protein having a transmembrane domain and of a recombinant of the human cDNA.		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

Internat'l Application No  
PCT/JP 99/02226

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/12 C07K14/47 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HILLIER L. ET AL.: "WashU-NCI human EST Project." EMBL DATABASE ENTRY AA680184; ACCESSION NUMBER AA680184, 4 December 1997 (1997-12-04), XP002113993 abstract	1-6
A	YOKOYAMA-KOBAYASHI M. ET AL.: "A signal sequence detection system using secreted protease activity as an indicator." GENE, vol. 163, 1995, pages 193-196, XP002046435 cited in the application the whole document	1-6

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

1 September 1999

Date of mailing of the international search report

17. 12. 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B

# INTERNATIONAL SEARCH REPORT

Intern. Application No  
PCT/JP 99/02226

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KYTE J. ET AL.: "A SIMPLE METHOD FOR DISPLAYING THE HYDROPATHIC CHARACTER OF A PROTEIN" JOURNAL OF MOLECULAR BIOLOGY, vol. 157, no. 1, 5 May 1982 (1982-05-05), pages 105-132, XP000609692 ISSN: 0022-2836 cited in the application the whole document ---	1-6
A	MORI K. ET AL.: "Kidney-specific expression of a novel mouse organic cation transporter-like protein." FEBS LETTERS, vol. 417, 1997, pages 371-374, XP002113977 the whole document ---	1-6
A	SCHÖMIG E. ET AL.: "Molecular cloning and characterization of two novel transport proteins from rat kidney." FEBS LETTERS, vol. 425, 20 March 1998 (1998-03-20), pages 79-86, XP002113994 the whole document ---	1-6
A	TASHIRO K. ET AL.: "Signal sequence trap: A cloning strategy for secreted proteins and type I membrane proteins." SCIENCE, vol. 261, 1993, pages 600-603, XP000673204 the whole document ---	1-6
P,A	YOKOYAMA-KOBAYASHI M. ET AL.: "Selection of cDNAs encoding putative type II membrane proteins on the cell surface from a human full-length cDNA bank." GENE, vol. 228, no. 1-2, 4 March 1999 (1999-03-04), pages 161-167, XP002113979 the whole document -----	1-6

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/JP 99/02226

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

See extra sheet, Invention 1.

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/JP 99/02226

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.1, DNA encoding it, cDNA comprising SEQ.ID.10 or SEQ.ID.19, expression vector and transformed cell capable of expressing said DNA or cDNA.

2. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.2, DNA encoding it, cDNA comprising SEQ.ID.11 or SEQ.ID.21, expression vector and transformed cell capable of expressing said DNA or cDNA.

3. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.3, DNA encoding it, cDNA comprising SEQ.ID.12 or SEQ.ID.23, expression vector and transformed cell capable of expressing said DNA or cDNA.

4. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.4, DNA encoding it, cDNA comprising SEQ.ID.13 or SEQ.ID.25, expression vector and transformed cell capable of expressing said DNA or cDNA.

5. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.5, DNA encoding it, cDNA comprising SEQ.ID.14 or SEQ.ID.27, expression vector and transformed cell capable of expressing said DNA or cDNA.

6. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.6, DNA encoding it, cDNA comprising SEQ.ID.15 or SEQ.ID.29, expression vector and transformed cell capable of expressing said DNA or cDNA.

7. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.7, DNA encoding it, cDNA comprising SEQ.ID.16 or SEQ.ID.31, expression vector and transformed cell capable of expressing said DNA or cDNA.

8. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.8, DNA encoding it, cDNA comprising SEQ.ID.17 or SEQ.ID.33, expression vector and transformed cell capable of expressing said DNA or cDNA.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/JP 99/02226

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## 9. Claims: 1-6 (PARTIALLY)

Protein comprising SEQ.ID.9, DNA encoding it, cDNA  
comprising SEQ.ID.18 or SEQ.ID.35, expression vector and  
transformed cell capable of expressing said DNA or cDNA.